

JOURNAL OF AVIAN BIOLOGY

Research article

Flying stimulates the antioxidant system and protects against oxidative damage in a migratory songbird, yet diet quality has little effect

Kristen J. DeMoranville¹, Wales Carter¹, Clara Cooper-Mullin¹, Liam Corcoran¹, Barbara J. Pierce² and Scott R. McWilliams¹✉

¹University of Rhode Island, Kingston, RI, USA

²Sacred Heart University, Fairfield, CT, USA

Correspondence: Scott R. McWilliams (srmcwilliams@uri.edu)

Journal of Avian Biology

2025: e03379

doi: [10.1002/jav.03379](https://doi.org/10.1002/jav.03379)

Subject Editor: Barbara Helm
Editor-in-Chief: Staffan Bensch
Accepted 07 March 2025



Ecologically relevant factors such as exercise and diet quality can directly influence how multifaceted physiological systems work; however, little is known about how such factors directly and interactively affect key components of the antioxidant system in multiple tissues of migratory songbirds. We tested 3 main hypotheses across three tissues in European starlings fed diets with more or less antioxidants (anthocyanins) and long-chain omega-6 polyunsaturated fats (18:2n6) while being flight-trained in a wind tunnel. *Stimulatory effect of flight*: flight-training stimulated the antioxidant system in that 1) plasma oxidative damage (dROMs) was reduced during a given acute flight, and contrary to our predictions, 2) antioxidant capacity (OXY or ORAC) and oxidative damage in plasma (dROMs), flight-muscle, and liver (LPO) of flight-trained birds were similar to that of untrained birds (i.e. not flown in a wind tunnel). Flight-trained birds that expended more energy per unit time (kJ min^{-1}) during their longest, final flight decreased antioxidant capacity (OXY) the most during the final flight. *Dietary fat quality effect*: contrary to our predictions, dietary 18:2n-6 did not influence oxidative status even after flight training. *Dietary antioxidant effect*: flight-trained birds supplemented with dietary anthocyanins did not have higher antioxidant capacity in plasma (OXY), or liver and flight-muscle (ORAC) compared to untrained birds. Counterintuitively, oxidative damage (dROMs) was higher in flight-trained supplemented birds compared to unsupplemented birds after an acute flight. In sum, the antioxidant system of songbirds flexibly responded to changes in availability of dietary antioxidants as well as increased flight time and effort, and such condition-dependent, individual-level, tissue-specific responses to the oxidative costs of long-duration flights apparently requires recovery periods for maintaining oxidative balance during migration.

Keywords: antioxidant capacity, antioxidants, dietary fat quality, flight training, migration, oxidative damage



www.avianbiology.org

© 2025 The Author(s). Journal of Avian Biology published by John Wiley & Sons Ltd on behalf of Nordic Society Oikos

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Introduction

The challenges of maintaining oxidative balance for migratory birds


The multifaceted antioxidant system available to birds consists of macromolecular sacrificial molecules, endogenous enzymes, and dietary antioxidants that can mitigate damage by scavenging reactive species (Fig. 1A). These antioxidant defenses exist in all metabolically active tissues of an organism, and the liver, flight-muscle or pectoralis, and blood are of particular importance in migratory birds. The liver is a crucial food processing organ, especially for exercising birds that rely on fat to fuel flight (McWilliams et al. 2004, Guglielmo 2018) and the pectoralis is the major skeletal muscle used to power flight (Biewener 2011). The blood is metabolically active in birds (unlike in mammals) and produces reactive species (Stier et al. 2013). Measuring indices of circulating oxidative status is important since an unbalanced oxidative state in the plasma would indicate damage to crucial molecules transported by the plasma (e.g. fatty acids, hemoglobin, cytokines) to muscles and organs. Thus, these three tissues relied on by flying birds likely have high antioxidant capacities to protect against oxidative damage, yet it remains unknown how antioxidant capacity and oxidative damage of these tissues respond to flight training, dietary fats, and dietary antioxidants or how the different components of the multifaceted antioxidant system work together to protect individuals against oxidative damage.

During energetically challenging periods, all aerobic organisms must contend with an increased production of reactive species by either neutralizing the reactive species with a toolkit of antioxidants to minimize resulting oxidative damage (outlined in Fig. 1A), or damage must be restored using repair mechanisms (Halliwell and Gutteridge 2007, Costantini 2014, 2019). Long-distance flight increases metabolism (Swanson 2010, Corder and Schaeffer 2015, DeMoranville et al. 2019, Bishop and Guglielmo 2022) and poses a potential oxidative challenge for flying animals (McWilliams et al. 2021, 2022). In Fig. 1B, a simplified diagram of the vertebrate antioxidant system shows how the extent of the antioxidant system can change in response to exercise, and the associated increase in metabolism and reactive species production, and thus affect the level of damage to lipids, proteins, and DNA. For example, Nathusius' bats *Pipistrellus nathusii* that were captured and sampled during a migratory flight had higher circulating oxidative damage markers compared to individuals that were captured and rested for 18–24 hours (Costantini et al. 2018). Similarly, circulating protein damage and the antioxidant enzyme glutathione peroxidase (GPx) were higher in European robins *Erithacus rubecula* during a nocturnal migratory flight compared to resting individuals caught during the day, indicating that damage to muscle occurs with flight and the antioxidant system can respond rapidly (Jenni-Eiermann et al. 2014). These acute effects of flight were also demonstrated in

captive yellow-rumped warblers *Setophaga coronata coronata*, as protein carbonyls and superoxide dismutase (SOD) activity were higher in the pectoralis muscle immediately after a flight in a wind tunnel compared to individuals at rest (Dick and Guglielmo 2019). Similarly, flight-trained zebra finches *Taeniopygia guttata* had increased circulating oxidative damage compared to untrained sedentary individuals (Skrip et al. 2016). These studies together demonstrate that migratory bats and birds respond to such oxidative challenges by increasing antioxidant enzyme activities, depleting or augmenting non-enzymatic antioxidant capacity, and in these cases increasing oxidative damage. They also reveal that too little is known about tissue-specific oxidative status of migratory birds (Dick and Guglielmo 2019) and the extent to which an individual's circulating oxidative status reflects the oxidative state of muscles and organs at a given time (Costantini 2019, Frawley et al. 2021a). Additionally, measurements reflecting different levels of the Central Dogma (e.g. gene expression versus enzyme activities) and when measurements are collected in relation to flight (e.g. during flight or after recovery) influence the comparisons between circulating and tissue oxidative status. For example, in a companion study the enzyme activities of CAT in the liver and pectoralis and GPx in the liver were higher in European starlings *Sturnus vulgaris* after recovering from flight training compared to birds that were not flight trained, yet these activities did not correlate with the higher gene expression levels of these enzymes in trained birds compared to untrained birds (DeMoranville et al. 2022). Consistent with these results, Frawley et al. (2021a) employed a similar experimental design and observed lower GPx activity in the heart and SOD activity in the liver of trained European starlings compared to starlings not trained in a wind tunnel during recovery, indicating that research on how antioxidant capacity recovers after flight should be mechanistically studied (McWilliams et al. 2021). What remains to be determined for migratory songbirds, and is the focus of this study, is how the oxidative status of the flight-muscle, liver, and primarily plasma responds to both flight training and ecologically relevant differences in diet quality (i.e. antioxidants and fat composition).

Certain dietary fats challenge the antioxidant system. Birds rely primarily on fatty acids to fuel flight (McWilliams et al. 2004, Guglielmo 2018), and several species of migratory songbirds increase the amounts of unsaturated fats and especially long-chain polyunsaturated fats (PUFA) in their diet, fat stores, and in circulation during migration compared to non-migration periods (Pierce et al. 2004, Pierce and McWilliams 2005, Price et al. 2008, Smith and McWilliams 2010, Jensen et al. 2020, McWilliams et al. 2022). The potential benefits of consuming certain long-chain PUFA (e.g. 18:2n-6 PUFA) including enhanced efficiency of energy metabolism during exercise and reduced flight costs (Pierce and McWilliams 2014, Guglielmo 2018), are also associated with potential oxidative costs because PUFA are highly susceptible to oxidative damage (Halliwell and Gutteridge 2007, Skrip and McWilliams 2016, McWilliams et al. 2020).

(A) The multifaceted antioxidant system



(B) How the antioxidant system may respond to exercise and diet quality




Figure 1. (A) The multifaceted antioxidant system available to birds that consists of macromolecular sacrificial molecules, endogenous enzymes, and dietary antioxidants that can mitigate damage by scavenging reactive species (Cooper-Mullin and McWilliams 2016). Reactive species produced during and after flight (thick arrows and lines, left) can act as beneficial signaling molecules, but when reactive species are in excess and not quenched by the multifaceted antioxidant system (middle boxes) the reactive species can cause damage to lipids, proteins, and DNA (thin arrows and lines, right). The three general components of the antioxidant system shown include sacrificial molecules produced during normal metabolism (e.g. uric acid produced during protein catabolism in birds), upregulation of endogenous enzymes (e.g. superoxide dismutase (SOD), catalase (CAT), and glutathione peroxidase (GPx), as well as antioxidants consumed in the diet (e.g.

Figure 1. Continued.

water-soluble polyphenols such as anthocyanins or fat-soluble vitamin E). In this study, we measured in blood and tissues the non-enzymatic antioxidant capacity (OXY and ORAC, respectively) as well as three primary endogenous enzymes (as reported in DeMoranville et al. 2022) and the level of damage (d-ROMs and lipid peroxides, respectively) while manipulating the amount of flight training, dietary anthocyanins, and fat quality to determine their effects on these components of the antioxidant system in a migratory songbird. (B) A simplified diagram of the vertebrate antioxidant system (Skrip and McWilliams 2016) showing how the extent of the antioxidant system can change in response to exercise, and the associated increase in metabolism and reactive species production, and thus affect the level of damage to lipids, proteins, and DNA. In the current study, we experimentally manipulated (gray boxes) the amount of flight training (i.e. flight-trained in a wind tunnel for 15 days or not) and the diet (high or low polyunsaturated fats (PUFA) and anthocyanins) in order to test three hypotheses related to how the antioxidant system of birds flexibly responds to such manipulations. Flight training increases reactive species and so stimulates the non-enzymatic antioxidant components (i.e. OXY in plasma, ORAC in liver and pectoral muscle), non-enzymatic antioxidant capacity can be augmented with dietary antioxidants as well as non-dietary endogenous sources (e.g. uric acid, glutathione, estrogen, melatonin) (Cooper-Mullin and McWilliams 2016, Skrip and McWilliams 2016), thereby building antioxidant capacity compared to untrained birds (H1). Birds fed diets composed of more 18:2n-6 PUFA are more susceptible to oxidative damage and thus preventatively increase antioxidant capacity (H2). Birds fed dietary anthocyanins will increase non-enzymatic antioxidant capacity compared to birds not fed anthocyanins (H3). We predicted that flight-trained birds would maintain low levels of oxidative damage (d-ROMs in plasma, lipid peroxides in tissues) because of their increased antioxidant capacity.

The resulting lipid radicals often cause a self-perpetuating chain reaction damaging nearby PUFA and other molecules (Halliwell and Gutteridge 2007, Skrip and McWilliams 2016, Cooper-Mullin et al. 2019). Birds preferentially consuming 18:2n-6 PUFA during migration to enhance their metabolism (Pierce and McWilliams 2014) likely require an augmented antioxidant system to protect against the oxidative challenge that such PUFA pose (McWilliams et al. 2021), as visually depicted in Fig. 1B.

Dietary antioxidants augment the antioxidant system. Many songbird species select fruits that are high in antioxidants and fat during fall (post-breeding) migration, suggesting that antioxidant consumption may be important to protect against oxidative damage during this life history stage (Alan and McWilliams 2013, Bolser et al. 2013) and we have visually depicted this scenario in Fig. 1B. Water-soluble antioxidants such as anthocyanins are preferentially consumed by certain songbird species during fall migration and in cafeteria-style choice experiments (Schaefer et al. 2008, Alan et al. 2013, Bolser et al. 2013). Anthocyanins are potent antioxidants (Halliwell and Gutteridge 2007). Anthocyanins can directly or indirectly affect the antioxidant system and also have been shown to protect birds from an immune challenge (Catoni et al. 2008) and protect against some of the metabolic costs associated with flight training (Casagrande et al. 2020) as well as courtship (Carbeck et al. 2018) and reproduction in spring (Frawley et al. 2021b). These energy savings associated with consuming anthocyanins may allow birds to invest in increasing enzymatic or non-enzymatic antioxidants to prevent damage (McWilliams et al. 2020). Non-enzymatic antioxidant capacity increases with fat stores in songbirds on stopovers during migration (Costantini et al. 2007, Skrip et al. 2015) and is greatest during the pre-migratory fueling stage in shorebirds (Gutiérrez et al. 2019). Thus, hyperphagia associated with preparation for migration in birds may augment their antioxidant system by increasing fat-soluble antioxidants stored within newly accumulated fat stores or by increasing water-soluble antioxidant metabolites continuously released by gut microbes (Dogan Comert and Gokman 2017).

How does flight training, dietary fat, and dietary anthocyanins affect the oxidative status of the plasma and metabolic tissues in a migratory songbird?

To better understand the multifaceted antioxidant system of birds, we conducted a factorial experiment that manipulated three ecologically relevant factors (i.e. flight training, dietary fat, dietary antioxidants) to determine their effects on equivalent measures of lipid damage and antioxidant capacity in the plasma, liver, and flight muscle of a migratory songbird. These three experimental manipulations are visualized in Fig. 1B, and the figure legend relates our hypotheses described below to the visuals in the diagram.

We tested the following hypotheses:

1. *Stimulatory effect of flight* (H1): acute flight and flight-training stimulate the non-enzymatic antioxidant system and protect against oxidative damage in the plasma, liver, and pectoralis.
 - a. *Acute-effect of a long flight*: plasma antioxidant capacity decreases during a given long-duration flight and so enables circulating oxidative damage levels to remain low.
 - b. *Long-term effect of flight-training*: a bird's plasma non-enzymatic antioxidant capacity (OXY) increases and lipid damage (d-ROMs) decreases over the course of several weeks of daily flight training as compared at discrete timepoints during the experiment.
 - c. *Flight-training effect across multiple tissues*: flying regularly over several weeks increases non-enzymatic antioxidant capacity (oxygen radical absorbance capacity, ORAC in flight muscle and liver; OXY in plasma) and decreases lipid damage (lipid hydroperoxides, LPO, in flight muscle and liver; d-ROMs in plasma) in a consistent manner across tissues compared to untrained birds not flown in a wind tunnel.
2. *Dietary fat effect* (H2): migratory songbirds fed diets composed of more 18:2n-6 PUFA are more susceptible to oxidative damage and thus preventatively increase antioxidant capacity in the plasma, liver, and flight muscle so as

to maintain low levels of lipid damage compared to birds fed diets with less PUFA.

3. *Dietary antioxidant effect* (H3): migratory songbirds fed dietary anthocyanins have increased non-enzymatic antioxidant capacity, and lower levels of lipid damage in all three tissues (i.e. liver, muscle, plasma) compared to birds not fed anthocyanins.

We also examined whether these three ecologically relevant factors (flying, fat quality of diet, dietary antioxidants) interacted to affect key components of the antioxidant system.

Material and methods

Experimental design

Omnivorous migratory songbirds undergo endurance flights biannually and many species switch to eating mostly berries that are rich in fats and antioxidants during their fall (post-breeding) migration (Alan et al. 2013, Bolser et al. 2013). Thus, they are an ideal natural system to study how the endogenous antioxidant system responds to flight training, dietary antioxidants, and dietary fat. We used European starlings *Sturnus vulgaris* as representative songbirds for this study because they are abundant in the New World and Old World, are omnivorous, acclimate well to captivity and new diets, and have been successfully trained and flown in wind tunnels in other studies (Nebel et al. 2012, Hall et al. 2014, Casagrande et al. 2020, McWilliams et al. 2020). Hatch year European starlings were caught at a dairy farm 20 km north of the Advanced Facility for Avian Research (AFAR), University of Western Ontario, London, Ontario, prior to fall migration in 2015 (19–23 August). European starlings in the Old World migrate from many hundreds of kilometers to several thousand kilometers (Feare 1984) mostly during daylight (Gruys-Casimir 1965, Feare 1984), although some migrate at night (Vigants et al. 2023). European starlings were introduced to North America in 1890–1891 and are now distributed across most of the continent with populations inhabiting the Great Lakes region, including Ontario, Canada, remaining migratory even though sedentary populations occur in other regions (Cabe 2020). Starlings were housed in one of four large indoor aviaries at the Advanced Facility for Avian Research (two 2.4 × 3.7 × 3.1 m and two 2.4 × 2.3 × 3.5 m). On 24 August we measured morphological characteristics, body mass, and molt score (0–5; Ginn and Melville 1983) for each individual. Birds were then randomly sorted into four groups (Fig. 2) with roughly equal distributions of body size and molt score. We maintained aviaries at 21°C on a natural light cycle from capture until the start of the experiment on 21 September when we fixed the light schedule at 11:13 L:D (day length on this date in London, Ontario). Upon capture and until the start of the experiment each week we weighed and inspected all birds to assess their health. All birds were cared for under animal care protocols for University of Western Ontario (2010–216) and the University of Rhode Island (AN11-12-009).

Experimental diets

Birds had ad libitum access to one of two semi-synthetic diets that had the same macronutrient content and differed only in fatty acid composition. The macronutrient composition of the two semisynthetic diets (41% carbohydrate, 13% protein, 30% fat) simulates a natural high-lipid fruit diet (Johnson et al. 1985, Smith et al. 2007), and the primary fatty acids in the diet (> 90% 16:0, 18:1, and 18:2) are also the most common fatty acids in natural fruits (Pierce and McWilliams 2014) and in songbirds that eat fruits during their migration (Pierce and McWilliams 2005, 2014). We manipulated the proportions of canola, sunflower, and palm oil so that the diets were either high (32%) or low (13%) in 18:2n-6 PUFA (omega-6 fatty acid, linoleic acid) which was primarily traded off with 16:0 (saturated fatty acid, palmitic acid). Thus, our experimental design requires us to attribute any observed dietary fat effects to both 18:2n-6 and 16:0 content. However, our interpretations focus on the potential effects of 18:2n-6 due to its demonstrated importance in metabolic signaling (Forman et al. 1997, Kennedy et al. 2007, Hamilton et al. 2018, Dick and Guglielmo 2019, Price et al. 2022). The complete list of diet ingredients and amounts have been previously published (Carter et al. 2020). Starlings in two aviaries received a 13% 18:2n-6 diet and two others received a 32% 18:2n-6 diet. The two diets have been shown to produce reliable differences in tissue fatty acid composition of starlings (Carter et al. 2020). On 1 September, after we were confident birds were well acclimated to the fat quality in the semisynthetic diets, we began adding a supplementary water-soluble antioxidant, anthocyanin (elderberry powder; Artemis International, Fort Wayne, IN) to the diets of birds in one 13% 18:2n-6 aviary and one 32% 18:2n-6 aviary, producing a 2 × 2 factorial diet manipulation with four diet groups (Fig. 2): 13% 18:2n-6, anthocyanin unsupplemented (n = 23), 13% 18:2n-6, anthocyanin supplemented (n = 23), 32% 18:2n-6, anthocyanin unsupplemented (n = 21), and 32% 18:2n-6, anthocyanin supplemented (n = 20). We chose the anthocyanin concentration used by researchers studying the effects of anthocyanin supplementation on food choice and immunocompetence in European blackcaps *Sylvia atricapilla* (Catoni et al. 2008, Schaefer et al. 2008). The anthocyanin supplement was equal to eating 2.8 mg per day which is equal to consuming 17 berries per day based on an average daily synthetic diet consumption of ca 35 wet g day⁻¹ (as observed in food intake trials in this study).

Experimental timeline

On 21 September we randomly assigned groups of 5 starlings to each of twenty cohorts (Fig. 2) although this cohort selection was stratified by extent of pre-basic I or post-juvenile molt (Ginn and Melville 1983) – birds furthest along in molt were flight-trained earliest. This stratified random sampling of individuals ensured that all birds had completed their flight feather molt prior to flight training. One cohort from each of the four diet groups was tested sequentially over 10




Figure 2. Experiment timeline included an initial acclimation to diets containing either 13% 18:2n-6 or 32% 18:2n-6 and then the addition of an antioxidant supplement to the diets of half the birds on 1 September to produce the 2 × 2 factorial diet manipulation with four diet groups: 13% 18:2n-6, anthocyanin unsupplemented (n = 23), 13% 18:2n-6, anthocyanin supplemented (n = 23), 32% 18:2n-6, anthocyanin unsupplemented (n = 21), and 32% 18:2n-6, anthocyanin supplemented (n = 20). After at least one month of acclimation to one of these four diets, birds were assigned to one of 20 cohorts each comprising 3 flight-trained birds and 2 untrained birds from each diet treatment (5 cohorts per treatment). On 23 September, and continuing every three days thereafter, a cohort began the four days of pre-training (shown by the white dotted box labelled PT) followed by fifteen days of flight training (shown by the dark box with vertical lines labelled FT). The 25-day flight-training schedule ended with birds performing a long-duration flight (pink shaded) followed by two recovery days (gray shaded). Blood sampling timepoints are indicated by the blood droplets and include for all birds a Background (BG) blood sample and a Recovery (RC) blood sample, and for flight-trained birds a Pre-flight (PF) and After flight (AF) blood sample taken the morning 1 day before the long-duration flight and immediately after the long-duration flight. Liver and pectoralis samples were taken from all birds on the final day. The acute effects of flight were assessed by comparing the change in plasma OXY and d-ROMs at the After-flight and Pre-flight time points. The long-term effects of flight-training on the antioxidant system were assessed two ways: (a) by comparing the change in plasma OXY and d-ROMs at the Recovery and Background time points, and (b) by comparing the oxidative status in three tissues (blood, liver, pectoralis) of flight-trained and untrained birds at the Recovery time point.

days, although the sampling order of the diet groups was randomly assigned so a diet group was not consistently sampled first or last. Thereafter another set of cohorts from each of the four diet groups was tested and so forth until the 20 cohorts of starlings had been tested by early December.

On 23 September, and continuing every three days, the 5 individuals from each selected cohort were removed from their aviaries, and we randomly assigned 2 birds as untrained birds (that do not undergo any flight-training) and 3 birds as flight-trained birds. Each selected cohort was initially placed in individual cages (0.6 × 0.5 × 0.5 m) for two days (days -9 and -8) to measure food intake and another two days (days -7 and -5) to measure basal and peak metabolic rates (Carter et al. 2020). On day -5 we returned the two untrained birds to their original aviary and moved the 3 flight-trained birds to a 0.8 × 1.5 × 2 m flight aviary.

Each cohort was blood sampled at consistent time points throughout the 25–26 day experimental period (Fig. 2: Background, BG; Pre-flight, PF; After-flight, AF; Recovery, RC). Birds were fasted for at least 1 hour before all blood sampling time points and were bled within 30 min of capture. Blood samples were taken from the brachial vein, and within 10 min of blood sampling the plasma was separated from the red blood cells following centrifugation at 11 000

g for 10 min (Damon/IEC Division, IEC MB centrifuge, micro hematocrit). Plasma was stored at -80°C until OXY-adsorbent test and d-ROM analyses. A blood sample was obtained 9 days prior to the start of flight-training at 08:00 h to obtain background oxidative measurements (Background) for each individual.

Flight training

In order to assess the impact of diet and endurance flight on the endogenous antioxidant system, three flight-trained birds were flown in a wind tunnel for four days of pre-training followed by fifteen days of flight training. Such a flight training regime has demonstrated success at eliciting long-duration flights in starlings (Engel et al. 2006). The wind tunnel was set to 12 m s⁻¹ windspeed, 15°C, and 70% humidity, and birds were fasted for 1 hour prior to all flights. Pre-training (PT; days -4 to -1) consisted of allowing training birds to fly between their flight cage and the wind tunnel followed by 20 min of habituation time per day in the wind tunnel with a perch. These initial four 'pre-training' days were not included in the reported overall training time, since birds could rest when needed. Starlings in the flight-training group then participated in a fifteen-day training regimen (flight

training, FT) that was used to test the long-term effects of antioxidant and damage measures among flight-trained and untrained birds (H1b). The flight-training regime consisted of increasing periods of flight (20–180 min) in the wind tunnel as follows: days 1–4, 20 min each day; day 5–6, 30 min each day; day 7, 60 min; day 8, 90 min; day 9, 30 min; day 10, 120 min; day 11, 180 min; day 12, rest day; day 13, 60 min; and day 14, 30 min. This flight training culminated in a flight on day 15 that lasted as long as birds would voluntarily fly, up to 6 h, and this flight was used to test the effects of acute flight on antioxidant and damage measures in the plasma of flight-trained birds (H1a). To determine fuel use during flight, body condition (fat and lean masses) was measured using a quantitative magnetic resonance machine (QMR; Echo Medical Systems, Houston, TX) immediately before and after the final flight. Energy expenditure during the flight was estimated by multiplying the mass of fat and lean tissue lost during flight by their respective energy densities, adding them, and dividing by flight duration (full methods and results reported previously (Carter et al. 2020)). The final flight was on average 193 ± 71 (SD) min and the maximum was 360 min. In order to test the acute effects of flight, we blood sampled flight-trained birds on the morning of day 14 at 08:00 h before their 30 min flight (Pre-flight, PF), this is at the same time that they would have been blood sampled before their longest flight on day 15. We sampled birds the morning prior to their long-flight to avoid excess stress associated with handling immediately prior to their long-flight. Since all experimental conditions were the same among the two days, we assume that this blood sample reflects the state of the flight-trained birds before a flight. An After-flight (AF) blood sample was taken immediately after the experimental flight (day 15) ranging from 11:00–14:00 h. Flight-trained birds were returned to their flight cages for two days to recover from their last flight. At 14:00–15:00 h on days 16 and 17 the untrained and trained birds, respectively, in each cohort were blood sampled for the final Recovery sample (RC). Birds were euthanized by cervical dislocation while under isoflurane anesthesia and the liver and pectoralis muscle samples were collected and immediately weighed. The liver and pectoralis were flash frozen in liquid nitrogen and stored at -80°C until analysis. This sampling design allowed us to compare oxidative status in the liver and pectoralis of untrained (control) birds and flight-trained birds that had recovered (for 48 h) from their longest flight on day 15.

Non-enzymatic antioxidant capacity and oxidative damage

OXY Adsorbent test. OXY was measured in the plasma (concentration unit: mmol l^{-1} of HClO neutralized; Diacron International, Grosseto, Italy). OXY directly measures the ability of a plasma sample to quench the oxidant hypochlorous acid. In addition to directly reacting with biological molecules, hypochlorous acid can form more deleterious hydroxyl radicals (Candeias et al. 1993). OXY provides an index of non-enzymatic antioxidant capacity (e.g. dietary

and non-dietary antioxidants), without being complicated by inclusion of uric acid (Costantini 2011, Alan and McWilliams 2013, Skrip and McWilliams 2016, Cooper-Mullin et al. 2019).

Oxygen radical absorbance capacity. In preparation for the measurement of oxygen radical absorbance capacity (ORAC) and lipid hydroperoxide concentration (LPO), approximately 250 mg of liver or pectoralis was homogenized on ice in 9 volumes of 0.1 M phosphate buffer, pH 7 with 3×10^5 s pulses of a high-speed stainless-steel homogenizer (Tissue Master 125, Omni International, Kennesaw GA USA). Homogenate was centrifuged at $10\,000\text{ g}$ for 10 min at 4°C (Beckman Coulter Allegra 21R), and the supernatant was aliquoted to two separate tubes to conduct the two separate assays (ORAC, LPO). Supernatant was immediately frozen at -80°C until the time of the assays (5–8 months after homogenization). We estimated antioxidant capacity against two of the more damaging forms of reactive oxygen species that readily react with lipids (Halliwell and Gutteridge 2007), peroxy and hydroxyl radicals (concentration unit: arbitrary units per gram of tissue, aug^{-1}), by using a microplate-based version of the competitive ORAC assay (Cao and Prior 1999) following Jimenez et al. 2020. When in vitro production of the radicals exceeds the antioxidant capacity of the tissue, these reactive oxygen species modify the algal pigment phycoerythrin (545/575 nm) and decrease its fluorescence. Peroxyl radicals were generated by 320 mmol l^{-1} 2,2'-azobis (2-amidinopropane) dihydrochloride, and hydroxyl radicals were generated in separate plates by adding $0.25\text{ }\mu\text{l}$ per well of 10 mmol l^{-1} CuSO_4 and 0.667 mol l^{-1} ascorbate mixture. ORAC values for peroxy and hydroxyl radicals were determined by integrating the area under the fluorescence decay curve.

Uric acid. Uric acid is a by-product of protein catabolism that acts as an antioxidant (Halliwell and Gutteridge 2007). Methods and results for this metabolite were reported previously (Carter et al. 2020). Briefly, we assayed uric acid concentration (concentration unit: mmol l^{-1}) using an absorbance endpoint assay adapted for small volumes (TECO Diagnostics, Anaheim, CA) on 96-well plates in duplicate.

Lipid oxidative damage d-ROMs. Oxidative damage in the plasma was measured using the d-ROMs test (concentration unit: mmol l^{-1} H_2O_2 equivalents; Diacron International). This test works by first decreasing the pH of the plasma to release metal ions from proteins to cleave circulating reactive oxygen metabolites (ROMs) through incubation with a solution of 0.01 mol l^{-1} acetic acid/sodium acetate buffer. The subsequent products react with a chromogen (N,N-diethyl-p-phenylenediamine) which has a color intensity that is proportional to the concentration of reactive oxygen metabolites in the plasma and was measured at 505 nm (Costantini et al. 2007, Costantini 2016, Cooper-Mullin et al. 2019). ROMs measured in this test are primarily hydroperoxides, and in plasma are primarily produced when reactive species interact with lipids (Davies 2016, Ito et al. 2017).

Lipid hydroperoxides. Oxidative damage in the liver and pectoralis was measured using the LPO test (concentration unit: hydroperoxide concentration μM ; Cayman Chemical,

Ann Arbor, MI). This assay provides a general measure of lipid peroxidation by directly measuring hydroperoxides compared to assays that measure the byproducts of specific fatty acid peroxidation (e.g. MDA, 4-HNE). Hydroperoxides in the sample were extracted using a chloroform and methanol method. Hydroperoxides in the extraction react with ferrous ions detected by thiocyanate ion chromogen and were read at 500 nm in a glass 96-well plate.

Plasma or tissue from each individual was measured in triplicate for LPO and in duplicate for all other assays (OXY, peroxy and hydroxyl radical absorbance capacity, d-ROMs); all coefficient of variations were under 10%, and replicates were averaged prior to statistical analyses.

Statistics

Model selection

We used R (ver. 3.5.3; www.r-project.org) for all analyses. We constructed either linear mixed effects models using the 'lme4' package or linear models using the 'stats' package to test our four main hypotheses. Final best-fit models were selected using an AIC selection criterion, $\Delta\text{AIC} \geq 3$. The only exception was when ΔAIC was within 3 for the models for liver and pectoralis lipid damage including cohort as a random factor compared with excluding it, and in these cases, we decided to control for cohort.

Linear mixed effects models

Linear mixed effects models including bird identification number and cohort number as random factors were used to analyze the longitudinal plasma data to test the hypotheses (H1a) that acute flight decreases antioxidant capacity during flight and enables oxidative damage levels to remain low, and (H1b) that regularly flying increases antioxidant capacity and oxidative damage. Linear mixed models that controlled for experimental cohort as a random factor were also used to test the hypothesis that flight-training (H1c), dietary fat (H2), and dietary antioxidants (H3) influenced change (indicated using Δ) during training of plasma lipid damage and lipid damage in the liver and pectoralis. We conducted pairwise comparisons using least square means to determine the differences in oxidative status between specific timepoints.

Linear models

Linear models were constructed to test the hypothesis that flight training (H1), dietary fat (H2), and dietary antioxidants (H3) influenced antioxidant capacity and lipid damage in the plasma at discrete timepoints (i.e. Background, Pre-flight, After-flight, Recovery), over the course of the experiment (calculated as Recovery–Background), during an acute flight (calculated as After-flight–Pre-flight), and in the liver and pectoralis at Recovery sampling. We used a global model without interaction terms that best matched this hypothesis and included possible explanatory covariates (i.e. cohort, sex, and wing chord); however, these covariates were not included in the global linear models.

To test the hypothesis that flight-training, dietary fat, and dietary antioxidants had an interactive effect on oxidative status, we compared our global models to models including a 3-way interaction between dietary fat, antioxidants, and training treatment. These models also tested the 2-way interactions between covariates. The models with the 3-way interactions were not among the best fit models, thus we report results for only the main effects.

To determine if energy expenditure during flight influenced antioxidant capacity or oxidative damage, we included energy consumed (kJ min^{-1}) during the acute flight as a fixed effect in linear models that compared both plasma oxidative status measured After-flight and the change in these measures during the acute flight (calculated as After-flight–Pre-flight) among diet groups across timepoints. The best fit models for After-flight oxidative status included energy expenditure, whereas those for change in oxidative status during the acute flight included energy expenditure \times diet interaction term.

Results

Stimulatory effect of flight training across time: acute-effect of a long flight (H1a)

Regular bouts of acute flight influenced OXY and d-ROMs in the plasma of flight-trained birds (i.e. comparison of Background, Pre-flight, After-flight, and Recovery blood samples; Fig. 3). Plasma d-ROMs was highest after two weeks of daily flight training compared to all other timepoints (Fig. 3; Background versus Pre-flight: $T_{49} = -3.042$, $p = 0.0027$, Pre-flight versus Recovery: $T_{49} = 3.085$, $p = 0.0023$). Immediately after an on average 193 ± 71 min flight the next day, plasma d-ROMs had substantially decreased relative to Pre-flight (Fig. 3A, Supporting information; Pre-flight versus After-flight: $T_{49} = 3.85$, $p = 0.0002$). In contrast, plasma OXY after two weeks of daily flight (Pre-flight) was substantially lower than Background and Recovery levels two days later (Fig. 3B, Supporting information; Background versus Pre-flight: $T_{49} = 6.058$, $p < 0.0001$, Pre-flight versus Recovery: $T_{49} = -6.113$, $p < 0.0001$). Similar to d-ROMs, immediately after a long-duration flight the next day, plasma OXY further decreased relative to Pre-flight (Fig. 3B, Supporting information; PF versus AF: $T_{49} = 2.330$, $p = 0.021$), and then returned to baseline levels at Recovery (Background versus After-flight: $T_{49} = 8.993$, $p < 0.0001$, After-flight versus Recovery: $T_{49} = -8.544$, $p < 0.0001$, Background versus Recovery: $T_{49} = 0.0694$, $p > 0.05$). These results support H1a (acute effects of flight) since plasma antioxidant capacity decreased during a given long-duration flight and levels of circulating oxidative damage remained lower than or at baseline levels.

Energy expenditure during flight implicated in regulating OXY depletion but not d-ROMs

Rate of energy expended during the flight (kJ min^{-1}) was not related to levels of plasma oxidative damage ($R^2 = 0.12$, Supporting information; Fig. 4A) but was related to the




Figure 3. Changes in (A) oxidative damage (d-ROMs) and (B) non-enzymatic antioxidant capacity (OXY) in the plasma (means \pm SD) in relation to flight-training. Background (BG) levels of d-ROMs and OXY (solid horizontal line \pm SD, dotted lines) were not significantly different for flight-trained and untrained birds (see text) and so were combined. Dietary antioxidants significantly affected only plasma d-ROMs measured After-flight (AF) so we discriminate between the two diets (filled or open triangles) for only this time point. Different lower-case letters for the 4 timepoints (Background BG, Pre-flight PF, After-flight AF, Recovery RC) for d-ROMs or OXY denote significant pairwise comparisons (Supporting information).

decrease in non-enzymatic antioxidant capacity during the acute flight ($R^2=0.39$, Supporting information; Fig. 4B). Only a quarter of birds experienced increased d-ROM levels during an acute flight, suggesting that most birds employed a successful antioxidant strategy to avoid the accumulation

of d-ROMs at all energy expenditures (Fig. 4A, note direction of arrow; Supporting information, note labels indicating d-ROM levels). Energy consumption during the flight did not explain After-flight levels of d-ROMs or the change in plasma d-ROMs during the flight (After-flight–Pre-flight;




Figure 4. The change in (A) oxidative damage (d-ROMs) and (B) non-enzymatic antioxidant capacity (OXY) in the plasma (means \pm SD) (After-flight–Pre-flight) related to the energy consumed during the flight (kJ min^{-1}). Birds generally decreased (inverted triangle) d-ROMs and OXY during flight; however, some birds increased (triangle) these measures. The individual variation in change in d-ROMs was not explained by energy consumed during flight. Whereas, birds that expended more energy had larger decreases in OXY however, this was driven by 7 birds that decreased OXY significantly more than the mean and consumed $> 0.51 \text{ kJ min}^{-1}$.

Fig. 4A, Supporting information; After-flight: $T_{46} = -2.018$, $p = 0.050$, Δ During flight d-ROMs: $T_{46} = -1.485$, $p = 0.149$). In contrast, OXY was lowest in birds that consumed more energy during flight (Supporting information; After-flight: $T_{46} = -3.133$, $p = 0.003$), and Δ During flight OXY decreased to the greatest extent in birds expending more energy (Fig. 4B, Supporting information; Δ During flight OXY: $T_{46} = -2.821$, $p = 0.007$). This relationship was primarily driven by seven individuals that decreased OXY by an average of -128 mmol l^{-1} of HClO neutralized (Fig. 4B and Supporting information), well below the mean decrease ($-22.7 \pm 10.4 \text{ mmol l}^{-1}$ of HClO neutralized). These birds expended energy in the upper 50% range ($> 0.51 \text{ kJ min}^{-1}$ was consumed by the upper 50% of individuals) (Fig. 4B and Supporting information), although other individuals that expended similar amounts of energy during flight did not have such extreme decreases in OXY. Instead, 5 of these birds increased uric acid by an average of 1.4 mmol , well above the mean increase ($0.5 \pm 0.06 \text{ mmol}$); whereas the remaining birds did not change these circulating measures (OXY, uric acid) from the averages.

Stimulatory effect of flight training across time: long-term effect of flight-training (H1b)

Circulating oxidative status (d-ROMs and OXY) was similar between untrained and flight-trained birds (Fig. 3A, Supporting information; Training: d-ROMs, $T_{279} = -0.400$, $p = 0.690$, OXY, $T_{279} = 0.623$, $p = 0.534$) at the start and end of the experiment (Background versus Recovery: d-ROMs,

$T_{90} = 0.054$, $p = 0.957$, OXY, $T_{90} = -0.069$, $p = 0.945$), suggesting that 15 days of flight training did not have substantial long-term effects on plasma oxidative status. However, the change in plasma d-ROMs over time (i.e. the difference from Recovery to Background) decreased more in flight-trained compared to untrained birds (Supporting information; Δ During training d-ROMs, $T_{89} = -2.025$, $p = 0.046$); we detected no such differences in change in OXY over time within individuals in relation to flight training (Δ During training OXY, $T_{89} = 0.982$, $p = 0.329$). These results provide partial evidence for H1b (Long-term effects of flying) as d-ROMs decreased over the course of several weeks of flying within flight-trained birds compared to untrained birds.

Stimulatory effect of flight-training across multiple tissues (H1c)

Lipid hydroperoxide concentration in the liver was lowest in flight-trained birds compared to untrained birds at Recovery (Fig. 5A; $T_{93} = 1.253$, $p < 0.001$). There were no differences in lipid hydroperoxide concentration in the pectoralis (Fig. 5A; $T_{95} = 1.000$, $p = 0.108$) or in plasma d-ROM levels (Supporting information; $T_{88} = 2.120$, $p = 0.107$) at Recovery between flight-trained and untrained birds. Antioxidant capacity was also not affected by 15 days of flight training in the plasma (Supporting information; Recovery, $T_{89} = 1.102$, $p = 0.274$), liver (Fig. 5B, Supporting information; Hydroxyl, $T_{92} = -0.744$, $p = 0.459$, Peroxyl, $T_{92} = -1.501$, $p = 0.137$), or pectoralis (Fig. 5B, Supporting information; Hydroxyl, $T_{92} = -1.939$, $p = 0.056$, Peroxyl, $T_{96} = 0.683$, $p = 0.496$).




Figure 5. (A) Lipid hydroperoxide concentration (LPO) and (B) Oxygen radical absorbance capacity (ORAC) (means \pm SD; Supporting information) in the liver and pectoralis muscle of European starlings that were or were not flown in the wind tunnel for 15 days, Flight-trained (n=49) or Untrained (n=40), respectively. Flight-trained birds had lower levels of lipid hydroperoxides in the liver compared to untrained birds but there were no differences in the pectoralis related to flight training. There were no significant differences in hydroxyl or peroxyl scavenging capacities between flight-trained and untrained birds in the liver or pectoralis. Asterisks correspond to significance levels: * $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$.

These lipid damage results provide evidence for the stimulatory effect of flight-training (H1c) in the liver but not in the pectoralis or plasma, whereas there is no evidence for such a stimulatory effect on antioxidant capacity.

Dietary fat (H2) and antioxidant (H3) effects on oxidative status

There were no effects of diet fat quality or dietary antioxidants on oxidative status of the liver or pectoralis (Supporting information), rather the effect of diet was limited to plasma d-ROMs and associated with acute flight (Fig. 3A, Supporting information). Contrary to hypothesis 3, d-ROM levels immediately after acute flight were lower in birds consuming diets unsupplemented with anthocyanins compared to birds supplemented with antioxidants (Supporting information; 13% PUFA, unsupplemented, $T_{46} = -2.161$, $p = 0.036$, 32% PUFA, unsupplemented, $T_{46} = -2.594$, $p = 0.013$, 32% PUFA, $T_{46} = -1.440$, $p = 0.586$), whereas anthocyanin supplementation did not affect OXY levels during acute flight (Supporting information; 13% PUFA, unsupplemented, $T_{46} = 0.492$, $p = 0.625$, 32% PUFA, unsupplemented, $T_{46} = -0.214$, $p = 0.832$, 32% PUFA supplemented, $T_{46} = -0.813$, $p = 0.421$). These results provide no support for H2 or H3 for all three tissues with the exception that dietary antioxidants influenced plasma lipid damage levels after flight.

Discussion

Acute- and long-term flight-stimulated antioxidant protection (H1)

We found evidence that repeated bouts of flight initiated a hormetic response that activated the antioxidant system to protect against the accumulation of oxidative damage, consistent with H1. This concept of hormesis, the mild exposure to reactive species and subsequent activation of protective and repair mechanisms, has been demonstrated mainly in the skeletal muscle of humans (McArdle et al. 2002, Rattan 2008). In the present study, the immediate, acute effects of a flight included a reduction in both non-enzymatic antioxidant capacity as well as oxidative damage (After-flight–Pre-flight; Fig. 3A, B), consistent with the acute-effects of flight hypothesis (H1a). The coincident reduction in both antioxidant capacity and oxidative damage suggests that birds effectively avoided oxidative damage caused by short-term exercise at least in part by using the non-enzymatic component of their antioxidant system. Similarly, birds exposed to 15 days of flight-training and two days of recovery were able to decrease circulating damage to lipids (Recovery–Background; Fig. 3A) while untrained birds were not (long-term flight effect, H1b). The apparent over-compensation of the antioxidant system to enable a decrease in oxidative damage during a given flight (H1a) seems novel to this study, as some flight-training studies report no change in markers of circulating damage after an acute flight (Skrip et al. 2016, Cooper-Mullin et al.

2019). Most studies report increases in markers of circulating oxidative damage shortly after a migratory flight in free-living birds (Jenni-Eiermann et al. 2014, Costantini et al. 2018), after a 200 km experimental flight in homing pigeons (Costantini et al. 2008), or in the pectoralis after an experimental flight in a wind tunnel (Dick and Guglielmo 2019). We found that the reduction in both non-enzymatic antioxidant capacity as well as oxidative damage was more apparent in birds that expended more energy per unit time during their longest flight while flying at a prescribed pace (Fig. 4). Perhaps flying birds that expend less energy per unit time are less metabolically challenged, and thus do not need to deplete OXY as much to prevent oxidative damage. This conclusion is similar in concept to Dick and Guglielmo 2019 who found less pectoralis damage in yellow-rumped warblers with lower flight energy expenditures.

Consistent with the flight effects proposed in H1b, regular daily flying depleted non-enzymatic antioxidant capacity with an associated decrease in oxidative damage immediately after a flight; although both non-enzymatic antioxidant capacity and oxidative damage returned to baseline levels after 48 h of rest (without wind tunnel flying). These results are in accordance with flight-trained zebra finches that also decreased OXY during 2 h of acute flight and then increased OXY after birds had a reprieve from regular flying (Cooper-Mullin et al. 2019). This consistent and relatively rapid (no more than a few days) depletion and recovery of non-enzymatic antioxidant capacity suggests that stopovers during migration may be important in allowing birds to maintain their oxidative status over the course of the entire migration (McWilliams et al. 2021). In support of this, plasma d-ROMs decreased with increasing stopover duration (0–8 nights) in garden warblers *Sylvia borin*, and plasma non-enzymatic antioxidant capacity increased with fat stores accumulated at a stopover site in blackpoll warblers *Setophaga striata* and red-eyed vireos *Vireo olivaceus* (Skrip et al. 2015). Importantly, after 15 days of exercise training, European starlings exposed to an acute flight further depleted their non-enzymatic antioxidant capacity without an associated increase in oxidative damage. This provides additional evidence that birds during migration seem able to rapidly adjust their antioxidant system to maintain overall low levels of circulating damage.

We also found evidence in the liver and pectoralis that flight training activated the antioxidant system so that after several weeks of daily flying the non-enzymatic antioxidant capacity was maintained at levels similar to those of untrained birds, and oxidative damage was prevented in all tissues, although this is contrary to our predictions that antioxidant capacity would be greater and oxidative damage levels would be lower in trained versus untrained birds (H1c). Repeated bouts of flight apparently protected birds against lipid damage in the liver and the pectoralis (Frawley et al. 2021a), as lipid hydroperoxide concentration was lower or similar in the liver and pectoralis, respectively, of flight-trained birds compared to untrained birds. In our companion study, DeMoranville et al. (2022), the gene expression of *CAT*, *SOD2*, and *GPXI* were upregulated in the liver of

flight-trained birds, while only *SOD2* was upregulated in the pectoralis, which likely explains how lower liver damage levels in flight-trained birds was achieved. This enzymatic upregulation combined with lower damage in the liver suggests that the liver is preferentially protected compared to the pectoralis, perhaps to preserve its crucial role in processing fatty acids during flight or to protect those fatty acids from degradation. Common quails *Coturnix coturnix* induced into a migratory phase using photoperiod had greater GPx activity in the pectoralis compared to birds not induced into a migratory phase, and this trend was detected in the liver of males but not females (Marasco et al. 2021). Liver damage, as measured by thiobarbituric acid reactive substances (TBARS), was higher in all birds in the migratory phase compared to the non-migratory phase, but TBARS in the pectoralis were similar among these groups, suggesting a potential preference to protect the pectoralis during the pre-migratory stage prior to migratory flights (Marasco et al. 2021). There were also no differences detected in plasma OXY or liver or pectoralis hydroxyl and peroxy scavenging capacities between flight-trained and untrained birds after 48 h of rest; thus birds were able to maintain constant long-term levels of antioxidant capacity when energetically challenged and avoid increases in oxidative damage. Although long-term antioxidant capacity in the tissues was unchanged, it is possible that flight-trained birds were able to prevent the accumulation of oxidative damage by utilizing non-enzymatic antioxidant capacity in tissues during flight, as also shown for OXY in the plasma, and by increasing the gene expression of enzymatic antioxidants (DeMoranville et al. 2022).

Dietary fat quality did not affect oxidative status (H2)

Our study does not provide evidence to support H2 that migratory songbirds fed diets composed of more 18:2n-6 are more susceptible to oxidative damage and thus must increase antioxidant capacity compared to when fed diets with less 18:2n-6. This hypothesis was informed by the biochemistry and oxidative susceptibility of PUFA (Halliwell and Gutteridge 2007, Skrip and McWilliams 2016) and the demonstrated responsiveness of the antioxidant system in migratory birds (Jenni-Eiermann et al. 2014, Skrip et al. 2015, Dick and Guglielmo 2019). Additionally, 18:2n-6 has been shown to stimulate antioxidant enzymes in fish (Li et al. 2013, Zengin and Yilmaz 2016) but not in rats (Tou et al. 2011). Dietary long-chain n-3 or n-6 PUFA did not affect oxidative damage or enzymatic antioxidants in the pectoralis of yellow-rumped warblers (Dick and Guglielmo 2019), and a companion study, DeMoranville et al. (2022), demonstrated that antioxidant gene expression and enzyme activities (GPx, SOD, CAT) did not increase in these same starlings consuming more 18:2n-6. The lack of change in antioxidant enzymes and oxidative damage among diets suggests that dietary 18:2n-6 composition did not oxidatively challenge birds even after flight training. It is possible that a bird's antioxidant system is equipped to combat reactive species associated with

consuming any amount of PUFA, which would explain the lack of differences among diet groups. For example, white-throated sparrows *Zonotrichia albicollis* fed diets with more 18:2n-3 and 18:2n-6 PUFA had higher levels of d-ROMs, but similar ratios of oxidative damage: antioxidant capacity (Alan and McWilliams 2013), and common blackbirds *Turdus merula* caught during migration had higher circulating total n-3 and n-6 PUFA and had higher non-enzymatic antioxidant capacity and similar damage levels compared to resident birds (Eikenaar et al. 2017, Jensen et al. 2020). Alternatively, common blackbirds may instead mediate total unsaturation levels in their diet and hence fat stores to minimize the oxidative challenge of consuming more PUFA (Jensen et al. 2020) rather than relying on their antioxidant system for protection. A companion study revealed that increased 18:2n-6 elevated the eicosanoid hormone prostacyclin which mediates and potentially increases the efficiency of lipid metabolism, and this could be another alternative explanation for the lack of an effect of dietary PUFA on the oxidative parameters measured in this study (d-ROMS, OXY, lipid hydroperoxides, ORAC) (Carter et al. 2024). In sum, lipid peroxidation likely remains a relevant challenge for migratory songbirds that rely on 18:2n-6 and other fats to fuel migratory flights (Pierce et al. 2004, Pierce and McWilliams 2005, Price et al. 2008, Smith and McWilliams 2010), and birds seem able to modulate their antioxidant system in response to the oxidative challenges fats pose and thereby successfully protect against damage. Whether or not these adjustments to the antioxidant system have metabolic tradeoffs or tradeoffs with immunity remains unknown (Costantini 2019). Thus, future studies that compare the effect of different amounts of dietary 18:2n-6 on a migratory bird's physiological status (e.g. oxidative, metabolic, and immune statuses) will better elucidate the direct effects of dietary 18:2n-6 on the potential tradeoffs among physiological systems.

Dietary antioxidants increased plasma lipid damage after flight (H3)

We also did not find support for H3 that migratory songbirds fed dietary anthocyanins have increased non-enzymatic antioxidant capacity, and lower levels of lipid damage in all three tissues compared to songbirds not fed anthocyanins. Anthocyanin supplementation did not affect non-enzymatic antioxidant capacity in the plasma or the two metabolic tissues. The same dietary anthocyanin concentration positively affected immune function in European black-caps (Caton et al. 2008, Schaefer et al. 2008), attenuated the corticosterone response to flight in European starlings (Casagrande et al. 2020), increased testosterone levels and breeding behaviors in male European starlings (Carbeck et al. 2018), and affected enzymatic antioxidant activity and lipid damage in the heart, non-enzymatic antioxidant capacity in the pectoralis, and protein damage in leg muscle (Frawley et al. 2021a). Thus, we were surprised to find that the same concentration of anthocyanins did not directly affect antioxidant capacity as measured in this study. Perhaps

when dietary anthocyanins are available, they are used in a tissue-specific manner for these other functions (immunity, metabolic protection) rather than directly for reactive species mitigation which then may allow any energy-cost savings to be invested in other aspects of the antioxidant system including, for example, enzymatic antioxidants or glutathione (the precursor to GPx) that are not detected in the OXY measurement. There is considerable cross-talk among inflammatory, immune, metabolic, and antioxidant pathways so that simultaneously measuring these multiple pathways seems necessary to understand potential trade-offs in response to dietary antioxidants (Costantini 2019).

Contrary to hypothesis 3, we found that d-ROM levels immediately after acute flight were lower in birds consuming diets without anthocyanins compared to birds supplemented with antioxidants. There are two potential explanations for this result. First, dietary anthocyanins prevented the accumulation of reactive species used to stimulate the endogenous antioxidant system, resulting in more oxidative damage. This preventative effect of consuming daily antioxidants (vitamin E and C in this case) decreased the expression of antioxidant enzymes and transcription factors in the skeletal muscle of humans (Ristow et al. 2009, Merry and Ristow 2016). Second, birds not consuming anthocyanins were more oxidatively challenged during flight which required them to upregulate endogenous antioxidants in red blood cells to reduce circulating damage. The circulating enzymatic antioxidant system can respond rapidly; for example, GPx activity increases in European robins during nocturnal migration and in zebra finches during experimentally imposed 2-h daily flights (Jenni-Eiermann et al. 2014, Cooper-Mullin et al. 2019), respectively. Given that water-soluble antioxidants such as anthocyanins are likely utilized as they are metabolized rather than stored for later use, as are lipid-soluble dietary antioxidants (Halliwell and Gutteridge 2007), perhaps this short half-life may be why we detected an antioxidant effect in the plasma in direct response to an acute oxidative challenge rather than in plasma or tissues after recovery from flight. We provide evidence that dietary anthocyanins influenced oxidative damage after an energetic challenge, yet to untangle the mechanistic underpinnings of this relationship, future studies that measure multiple antioxidant classes and evaluate the crosstalk among various physiological pathways are required.

Relevance and significance

The multi-faceted antioxidant system consists of numerous components that collectively provide individuals with a tool-box of 'antioxidant strategies' they can use to protect against oxidative damage (Halliwell and Gutteridge 2007, Costantini 2014). Migratory birds can use a combination of non-enzymatic antioxidants gained from their diets and a suite of endogenous antioxidant enzymes and sacrificial molecules to protect against excess reactive species produced by relevant oxidative challenges, including repeated bouts of

flight and high-PUFA diets (Fig. 1) (Skríp and McWilliams 2016, McWilliams et al. 2021). This study demonstrated the ability of a songbird to use OXY during an acute flight and to recover OXY levels within 2 days of rest (Fig. 3), revealing the recovery timeline of antioxidant capacity for the first time (McWilliams et al. 2021). We also found that such rapid adjustments to the antioxidant system allowed birds to maintain overall low levels of circulating damage, although the extent was tissue-dependent. Researchers should remain mindful of the tissue-dependency of antioxidant capacity and oxidative damage demonstrated here and elsewhere (Frawley et al. 2021a) and that future studies that assess antioxidant capacity and oxidative damage select biologically relevant tissues to measure that directly relate to the hypotheses of interest.

Intra-individual variation in the ability to utilize antioxidants in response to various energetically demanding scenarios must exist for phenotypic flexibility of the antioxidant system to persist (Piersma and Van Gils 2011) and we found evidence of such individual-level, condition-dependent modulation of OXY and uric acid during long-duration flight (Supporting information). A subset of birds that expended energy in the upper 50% range ($> 0.51 \text{ kJ min}^{-1}$) seemed to either deplete OXY or increase levels of uric acid, a potent antioxidant (Fig. 4B and Supporting information), whereas the remaining birds did not change these circulating measures (OXY, uric acid) from the averages. Studies, such as those done with mosquitofish (Loughland and Seebacher 2020), that elucidate the antioxidant strategies used by individuals to maintain their oxidative balance under oxidatively challenging conditions and its effects on performance (e.g. temperature acclimation, running speed, breeding success, flight efficiency) would be particularly revealing for migratory songbirds.

The study described here was part of a larger integrated research project that also compared metabolic and antioxidant gene expression and enzyme activities in multiple tissues within the same individuals (DeMoranville et al. 2020, 2022). This allows us to compare how flight training, dietary antioxidants, and dietary PUFA affect the antioxidant system across the traditional scales of the Central Dogma (DNA, to RNA, to protein). All measures (i.e. in the liver and pectoralis: select metabolic and antioxidant genes, antioxidant enzyme activities; in the plasma: OXY, d-ROMs) except lipid hydroperoxides and ORAC in tissues responded strongly to flight training. In the plasma, proteins are subject to post-translational modifications and so better able to detect individual strategies to combat oxidative damage in response to flight exercise. In contrast, diet effects on antioxidant capacity were most evident in the pectoralis and liver (i.e. relative gene expression and antioxidant assays) compared to functional assays in the plasma (i.e. markers for antioxidant capacity and oxidative damage), perhaps because the former is more tightly regulated. Our studies also suggest that plasma antioxidant capacity is not a good predictor of oxidative damage in tissues, whereas antioxidant gene expression and antioxidant enzyme activities predict oxidative damage in the

pectoralis and liver to varying extents as reported in a PCA analysis, although correlations were low among the antioxidant and damage measures (DeMoranville 2020). Thus, future studies must carefully consider the appropriate scale of measurement(s) to best reveal how the antioxidant system of wild animals responds to ecologically relevant oxidative challenges such as migration.

Acknowledgements – We would like to express gratitude to Artemis International for donating the elderberry extract used for the dietary anthocyanin supplementation. We appreciate Todd Hyland and the hospitality he offered at his Hyland Dairy Farm Ltd; thanks for letting us climb all over the rafters of your barn in pursuit of European starlings. We greatly appreciate Andrew Gould, Mikaela Rebuli, Chris Guglielmo, and the entire staff at AFAR for providing necessary infrastructure, animal care assistance, and logistical help throughout the experiment. Undergraduate research assistants, Luke Douglas and Kiera Bohannon, were an integral part of the research team and we appreciate their eagerness and exceptional work ethic. We would like to express our great appreciation to the McWilliams lab members, and Michael Weir's support throughout this dissertation was invaluable.

Funding – This research was funded by a National Science Foundation Grant, Award no. IOS:1354187, to SRM and BJP.

Permits – All birds were cared for under animal care protocols for University of Western Ontario (2010-216) and the University of Rhode Island (AN11-12-009). Samples brought from Canada to the USA were authorized by USDA (import permit no. 129139).

Author contributions

Kristen J. DeMoranville: Conceptualization (equal); Data curation (lead); Formal analysis (lead); Investigation (lead); Methodology (equal); Project administration (lead); Resources (equal); Software (equal); Supervision (lead); Validation (lead); Visualization (lead); Writing – original draft (lead); Writing – review and editing (lead). **Wales Carter:** Conceptualization (equal); Data curation (supporting); Investigation (equal); Methodology (equal); Project administration (supporting); Resources (equal); Supervision (supporting); Writing – review and editing (supporting). **Clara Cooper-Mullin:** Visualization (equal); Writing – review and editing (supporting). **Liam Corcoran:** Data curation (supporting); Writing – review and editing (supporting). **Barbara J. Pierce:** Conceptualization (lead); Funding acquisition (lead); Investigation (supporting); Methodology (equal); Project administration (supporting); Resources (equal); Supervision (equal); Writing – review and editing (supporting). **Scott R. McWilliams:** Conceptualization (lead); Funding acquisition (lead); Investigation (equal); Methodology (lead); Project administration (equal); Resources (equal); Supervision (lead); Writing – original draft (supporting); Writing – review and editing (equal).

Transparent peer review

The peer review history for this article is available at <https://www.webofscience.com/api/gateway/wos/peer-review/jv.03379>.

Data availability statement

Data are available from the Dryad Digital Repository: <https://doi.org/10.5061/dryad.dv41ns26w> (DeMoranville et al. 2025).

Supporting information

The Supporting information associated with this article is available with the online version.

References

- Alan, R. R. and McWilliams, S. R. 2013. Oxidative stress, circulating antioxidants, and dietary preferences in songbirds. – *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* 164: 185–193.
- Alan, R. R., McWilliams, S. R. and McGraw, K. J. 2013. The importance of antioxidants for avian fruit selection during autumn migration. – *Wilson J. Ornithol.* 125: 513–525.
- Biewener, A. A. 2011. Muscle function in avian flight: achieving power and control. – *Philos. Trans. R. Soc. B* 366: 1496–1506.
- Bishop, C. M. and Guglielmo, C. G. 2022. Flight. – In: Scanes, C. G. and Dridi, S. (eds), *Sturkie's avian physiology*, 7th edn. Academic Press, pp. 1265–1329.
- Bolser, J. A., Alan, R. R., Smith, A. D., Li, L., Seeram, N. P. and McWilliams, S. R. 2013. Birds select fruits with more anthocyanins and phenolic compounds during autumn migration. – *Wilson J. Ornithol.* 125: 97–108.
- Cabe, P. R. 2020. European starling (*Sturnus vulgaris*), ver. 1.0. – In: Billerman, S. M. (ed.), *Birds of the world*. Cornell Laboratory of Ornithology. <https://doi.org/10.2173/bow.eursta.01>.
- Cao, G. and Prior, R. L. 1999. Anthocyanins are detected in human plasma after oral administration of an elderberry extract. – *Clin. Chem.* 45: 574–576.
- Carbeck, K. M., DeMoranville, K. J., D'Amelio, P. B., Goymann, W., Trost, L., Pierce, B., Bryla, A., Dzialo, M., Bauchinger, U. and McWilliams, S. R. 2018. Environmental cues and dietary antioxidants affect breeding behavior and testosterone of male European starlings (*Sturnus vulgaris*). – *Horm. Behav.* 103: 36–44.
- Carter, W. A., DeMoranville, K. J., Pierce, B. J. and McWilliams, S. R. 2020. The effects of dietary linoleic acid and hydrophilic antioxidants on basal, peak, and sustained metabolism in flight-trained European starlings. – *Ecol. Evol.* 10: 1552–1566.
- Carter, W. A., DeMoranville, K. J., Trost, L., Bryla, A., Dzialo, M., Sadowska, E. T., Bauchinger, U., Pierce, B. and McWilliams, S. R. 2024. Dietary fatty acids and flight-training influence the expression of the eicosanoid hormone prostacyclin in songbirds. – *Comp. Biochem. Physiol. A* 288: 111561.
- Casagrande, S., DeMoranville, K. J., Trost, L., Pierce, B. J., Bryla, A., Dzialo, M., Sadowska, E. T., Bauchinger, U. and McWilliams, S. R. 2020. Dietary antioxidants attenuate the endocrine stress response during long-duration flight of a migratory bird. – *Proc. Natl Acad. Sci. USA* 18: 1–8.
- Catoni, C., Schaefer, H. M. and Peters, A. 2008. Fruit for health: the effect of flavonoids on humoral immune response and food selection in a frugivorous bird. – *Funct. Ecol.* 22: 649–654.
- Cooper-Mullin, C. and McWilliams, S. R. 2016. The role of the antioxidant system during intense endurance exercise: lessons from migrating birds. – *J. Exp. Biol.* 219: 3684–3695.

- Cooper-Mullin, C., Carter, W. A. and McWilliams, S. R. 2019. Acute effects of intense exercise on the antioxidant system in birds: does exercise training help? – *J. Exp. Biol.* 222: jeb210443.
- Corder, K. R. and Schaeffer, P. J. 2015. Summit metabolic rate exhibits phenotypic flexibility with migration, but not latitude in a neotropical migrant, *Parkesia noveboracensis*. – *J. Ornithol.* 156: 547–550.
- Costantini, D. 2014. Oxidative stress and hormesis in evolutionary ecology and physiology: a marriage between mechanistic and evolutionary approaches. – Springer-Verlag.
- Costantini, D. 2016. Oxidative stress ecology and the d-ROMs test: facts, misfacts and an appraisal of a decade's work. – *Behav. Ecol. Sociobiol.* 70: 809–820.
- Costantini, D. 2019. Understanding diversity in oxidative status and oxidative stress: the opportunities and challenges ahead. – *J. Exp. Biol.* 222: jeb194688.
- Costantini, D., Cardinale, M. and Carere, C. 2007. Oxidative damage and anti-oxidant capacity in two migratory bird species at a stop-over site. – *Comp. Biochem. Physiol. C* 144: 363–371.
- Costantini, D., Dell'Ariccia, G. and Lipp, H.-P. 2008. Long flights and age affect oxidative status of homing pigeons (*Columba livia*). – *J. Exp. Biol.* 211: 377–381.
- Costantini, D., Lindecke, O., Pétersons, G. and Voigt, C. C. 2018. Migratory flight imposes oxidative stress in bats. – *Curr. Zool.* 65: 147–153.
- Davies, M. J. 2016. Protein oxidation and peroxidation. – *Biochem. J.* 473: 805–825.
- DeMoranville, K. J. 2020. The frequent flyer's guide to migration: how songbird metabolism and oxidative status respond to endurance flight and diet quality. – Univ. of Rhode Island.
- DeMoranville, K. J., Corder, K. R., Hamilton, A., Russell, D. E., Huss, J. M. and Schaeffer, P. J. 2019. PPAR expression, muscle size and metabolic rates across the gray catbird's annual cycle are greatest in preparation for fall migration. – *J. Exp. Biol.* 222: jeb198028.
- DeMoranville, K. J., Carter, W. A., Pierce, B. J. and McWilliams, S. R. 2020. Flight training in a migratory bird drives metabolic gene expression in the flight muscle but not liver, and dietary fat quality influences select genes. – *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 319: R637–R652.
- DeMoranville, K. J., Carter, W. A., Pierce, B. J. and McWilliams, S. R. 2022. Flight and dietary antioxidants influence antioxidant expression and activity in a migratory bird. – *Integr. Org. Biol.* 4: obab035.
- DeMoranville, K. J., Carter, W., Cooper-Mullin, C., Corcoran, L., Pierce, B. J. and McWilliams, S. R. 2025. Data from: Flying stimulates the antioxidant system and protects against oxidative damage in a migratory songbird, yet diet quality has little effect. – Dryad Digital Repository, <https://doi.org/10.5061/dryad.dv41ns26w>.
- Dick, M. F. and Guglielmo, C. G. 2019. Flight muscle protein damage during endurance flight is related to energy expenditure but not dietary polyunsaturated fatty acids in a migratory bird. – *J. Exp. Biol.* 222: 1–9.
- Dogan Comert, E. and Gokman, V. 2017. Antioxidants bound to an insoluble food matrix: their analysis, regeneration behavior, and physiological importance. – *Compr. Rev. Food Sci. Saf.* 16: 382–339.
- Eikenaar, C., Källstig, E., Andersson, M. N., Herrera-Dueñas, A. and Isaksson, C. 2017. Oxidative challenges of avian migration: a comparative field study on a partial migrant. – *Physiol. Biochem. Zool.* 90: 223–229.
- Engel, S., Biebach, H. and Visser, G. H. 2006. Metabolic costs of avian flight in relation to flight velocity: a study in rose coloured starlings (*Sturnus roseus*, Linnaeus). – *J. Comp. Physiol. B* 176: 415–427.
- Feare, C. J. 1984. The starling. – Oxford Univ. Press.
- Forman, B. M., Chen, J. and Evans, R. M. 1997. Hypolipidemic drugs, polyunsaturated fatty acids, and eicosanoids are ligands for peroxisome proliferator-activated receptors α and δ . – *Proc. Natl Acad. Sci. USA* 94: 4312–4317.
- Frawley, A. E., DeMoranville, K. J., Carbeck, K. M., Trost, L., Bryła, A., Działo, M., Sadowska, E. T., Bauchinger, U., Pierce, B. J. and McWilliams, S. R. 2021a. Flight training and dietary antioxidants have mixed effects on the oxidative status of multiple tissues in a female migratory songbird. – *J. Exp. Biol.* 224: jeb243158.
- Frawley, A. E., DeMoranville, K. J., Carbeck, K. M., Trost, L., Bryła, A., Działo, M., Sadowska, E. T., Bauchinger, U., Pierce, B. J. and McWilliams, S. R. 2021b. Season, anthocyanin supplementation, and flight training have mixed effects on the antioxidant system of migratory European starlings. – *Ornithology* 138: 1–16.
- Ginn, H. and Melville, D. 1983. Molt in birds. – British Trust for Ornithology.
- Gruys-Casimir, E. M. 1965. On the influence of environmental factors on the autumn migration of chaffinch and starling: a field study. – *Arch. Néerl. Zool.* 16: 175–279.
- Guglielmo, C. G. 2018. Obese super athletes: fat-fueled migration in birds and bats. – *J. Exp. Biol.* 221: jeb165753.
- Gutiérrez, J. S., Sabat, P., Castañeda, L. E., Contreras, C., Navarrete, L., Peña-Villalobos, I. and Navedo, J. G. 2019. Oxidative status and metabolic profile in a long-lived bird preparing for extreme endurance migration. – *Sci. Rep.* 9: 17616.
- Hall, Z. J., Bauchinger, U., Gerson, A. R., Price, E. R., Langlois, L. A., Boyles, M., Pierce, B., McWilliams, S. R., Sherry, D. F. and Macdougall-Shackleton, S. A. 2014. Site-specific regulation of adult neurogenesis by dietary fatty acid content, vitamin E and flight exercise in European starlings. – *Eur. J. Neurosci.* 39: 875–882.
- Halliwell, B. and Gutteridge, J. M. 2007. Free radicals in biology and medicine. – Oxford Univ. Press.
- Hamilton, A., Ly, J., Robinson, J. R., Corder, K. R., DeMoranville, K. J., Schaeffer, P. J. and Huss, J. M. 2018. Conserved transcriptional activity and ligand responsiveness of avian PPARs: potential role in regulating lipid metabolism in migratory birds. – *Gen. Comp. Endocrinol.* 268: 110–120.
- Ito, F., Ito, T., Suzuki, C., Yahata, T., Ikeda, K. and Hamaoka, K. 2017. The application of a modified d-ROMs test for measurement of oxidative stress and oxidized high-density lipoprotein. – *Int. J. Mol. Sci.* 18: 454.
- Jenni-Eiermann, S., Jenni, L., Smith, S. and Costantini, D. 2014. Oxidative stress in endurance flight: an unconsidered factor in bird migration. – *PLoS One* 9: e97650.
- Jensen, J. K., Isaksson, C., Eikenaar, C. and Andersson, M. N. 2020. Migrant blackbirds, *Turdus merula*, have higher plasma levels of polyunsaturated fatty acids compared to residents, but not enhanced fatty acid unsaturation index. – *Ecol. Evol.* 10: 10196–10206.
- Jimenez, A. G., Ruhs, E. C., Tobin, K. J., Anderson, K. N., Le Pogam, A., Regimbald, L. and Vézina, F. 2020. Consequences of being phenotypically mismatched with the environment: no evidence of oxidative stress in cold- and warm-acclimated birds facing a cold spell. – *J. Exp. Biol.* 223: jeb218826.

- Johnson, R. A., Willson, M. F., Thompson, J. N. and Bertin, R. I. 1985. Nutritional values of wild fruits and consumption by migrant frugivorous birds. – *Ecology* 66: 819–827.
- Kennedy, S. R., Bickerdike, R., Berge, R. K., Porter, A. R. and Tocher, D. R. 2007. Influence of dietary conjugated linoleic acid (CLA) and tetradecylthioacetic acid (TTA) on growth, lipid composition and key enzymes of fatty acid oxidation in liver and muscle of Atlantic cod (*Gadus morhua* L.). – *Aquaculture* 264: 372–382.
- Li, M., Chen, L., Qin, J. G., Li, E., Yu, N. and Du, Z. 2013. Growth performance, antioxidant status and immune response in darkbarbel catfish *Pelteobagrus vachelli* fed different PUFA/vitamin E dietary levels and exposed to high or low ammonia. – *Aquaculture* 406–407: 18–27.
- Loughland, I. and Seebacher, F. 2020. Differences in oxidative status explain variation in thermal acclimation capacity between individual mosquitofish (*Gambusia holbrooki*). – *Funct. Ecol.* 34: 1380–1390.
- Marasco, V., Sebastiano, M., Costantini, D., Pola, G. and Fusani, L. 2021. Controlled expression of the migratory phenotype affects oxidative status in birds. – *J. Exp. Biol.* 224: jeb233486.
- McArdle, A., Vasilaki, A. and Jackson, M. 2002. Exercise and skeletal muscle ageing: cellular and molecular mechanisms. – *Ageing Res. Rev.* 1: 79–93.
- McWilliams, S. R., Guglielmo, C., Pierce, B. and Klaassen, M. 2004. Flying, fasting, and feeding in birds during migration: a nutritional and physiological ecology perspective. – *J. Avian Biol.* 35: 377–393.
- McWilliams, S. R., Pierce, B. J., Wittenzellner, A., Langlois, L., Engel, S., Speakman, J. R., Fatica, O., DeMoranville, K. J., Goymann, W., Trost, L., Bryla, A., Dzialo, M., Sadowska, E. T. and Bauchinger, U. 2020. The energy savings–oxidative cost trade-off for birds during migration. – *eLife* 9: e60626.
- McWilliams, S. R., Carter, W. A., Cooper-Mullin, C., DeMoranville, K. J., Frawley, A. E., Pierce, B. J. and Skrip, M. M. 2021. How birds during migration maintain (oxidative) balance. – *Front. Ecol. Evol.* 9: 1–17.
- McWilliams, S. R., Ramenofsky, M. and Pierce, B. J. 2022. Physiological challenges of migration. – In: Scanes, C. G. and Dridi, S. (eds), *Sturkie's avian physiology*, 2nd edn. Academic Press, pp. 1331–1372.
- Merry, T. L. and Ristow, M. 2016. Do antioxidant supplements interfere with skeletal muscle adaptation to exercise training? – *J. Physiol.* 594: 5135–5147.
- Nebel, S., Bauchinger, U., Buehler, D. M., Langlois, L. A., Boyles, M., Gerson, A. R., Price, E. R., McWilliams, S. R. and Guglielmo, C. G. 2012. Constitutive immune function in European starlings, *Sturnus vulgaris*, is decreased immediately after an endurance flight in a wind tunnel. – *J. Exp. Biol.* 215: 272–278.
- Pierce, B. J. and McWilliams, S. R. 2005. Seasonal changes in composition of lipid stores in migratory birds: causes and consequences. – *Condor* 107: 269–279.
- Pierce, B. J. and McWilliams, S. R. 2014. The fat of the matter: how dietary fatty acids can affect exercise performance. – *Integr. Comp. Biol.* 54: 903–912.
- Pierce, B. J., McWilliams, S. R., Place, A. R. and Huguenin, M. A. 2004. Diet preferences for specific fatty acids and their effect on composition of fat reserves in migratory red-eyed vireos (*Vireo olivaceus*). – *Comp. Biochem. Physiol. A* 138: 503–514.
- Piersma, T. and Van Gils, J. A. 2011. The flexible phenotype: a body-centred integration of ecology, physiology, and behaviour. – Oxford Univ. Press.
- Price, E. R., Krokfors, A. and Guglielmo, C. G. 2008. Selective mobilization of fatty acids from adipose tissue in migratory birds. – *J. Exp. Biol.* 211: 29–34.
- Price, E. R., Bauchinger, U., McWilliams, S. R., Boyles, M. L., Langlois, L. A., Gerson, A. R. and Guglielmo, C. G. 2022. The effects of training, acute exercise and dietary fatty acid composition on muscle lipid oxidative capacity in European starlings. – *J. Exp. Biol.* 225: jeb244433.
- Rattan, S. I. S. 2008. Hormesis in aging. – *Ageing Res. Rev.* 7: 63–78.
- Ristow, M., Zarse, K., Oberbach, A., Klötting, N., Birringer, M., Kiehntopf, M., Stumvoll, M., Kahn, C. R. and Blüher, M. 2009. Antioxidants prevent health-promoting effects of physical exercise in humans. – *Proc. Natl Acad. Sci. USA* 2: 2–7.
- Schaefer, H. M., McGraw, K. and Catoni, C. 2008. Birds use fruit colour as honest signal of dietary antioxidant rewards. – *Funct. Ecol.* 22: 303–310.
- Skrip, M. M. and McWilliams, S. R. 2016. Oxidative balance in birds: an atoms-to-organisms-to-ecology primer for ornithologists. – *J. Field Ornithol.* 87: 1–20.
- Skrip, M. M., Bauchinger, U., Goymann, W., Fusani, L., Cardinale, M., Alan, R. R. and McWilliams, S. R. 2015. Migrating songbirds on stopover prepare for, and recover from, oxidative challenges posed by long-distance flight. – *Ecol. Evol.* 5: 3198–3209.
- Skrip, M. M., Seeram, N. P., Yuan, T., Ma, H. and McWilliams, S. R. 2016. Dietary antioxidants and flight exercise in female birds affect allocation of nutrients to eggs: how carry-over effects work. – *J. Exp. Biol.* 219: 2716–2725.
- Smith, S. B. and McWilliams, S. R. 2010. Patterns of fuel use and storage in migrating passerines in relation to fruit resources at autumn stopover sites. – *Auk* 127: 108–118.
- Smith, S. B., McPherson, K. H., Backer, J. M., Pierce, B. J., Podlesak, D. W. and McWilliams, S. R. 2007. Fruit quality and consumption by songbirds during autumn migration. – *Wilson J. Ornithol.* 119: 419–428.
- Stier, A., Bize, P., Schull, Q., Zoll, J., Singh, F., Geny, B., Gros, F., Royer, C., Massemin, S. and Criscuolo, F. 2013. Avian erythrocytes have functional mitochondria, opening novel perspectives for birds as animal models in the study of ageing. – *Front. Zool.* 10: 33.
- Swanson, D. L. 2010. Seasonal metabolic variation in birds: functional and mechanistic correlates. – In: Thompson, C. (ed.), *Current ornithology*. Vol. 17. Springer, pp. 131–189.
- Tou, J. C., Altman, S. N., Gigliotti, J. C., Benedito, V. A. and Cordonier, E. L. 2011. Different sources of omega-3 polyunsaturated fatty acids affects apparent digestibility, tissue deposition, and tissue oxidative stability in growing female rats. – *Lipids Health Dis.* 10: 179.
- Vigants, V., Keišs, O., Dinsbergs, I., Jaunzemis, V., Zacmane, E. M., Priedniece, A. and Briedis, M. 2023. Migration strategies, performance and annual activity budget in a short-distance migrant, the common starling *Sturnus vulgaris*. – *J. Avian Biol.* 2023: e03080.
- Zengin, H. and Yilmaz, Ö. 2016. Antioxidant defence of the actively feeding *Oncorhynchus mykiss* (Walbaum 1792) larvae in relation to dietary PUFA and vitamin E contents. – *Reg. Stud. Mar. Sci.* 8: 515–522.