

Research



Cite this article: Slezak CR *et al.* 2024
Unconventional life history in a migratory
shorebird: desegregating reproduction and
migration. *Proc. R. Soc. B* **291**: 20240021.
<https://doi.org/10.1098/rspb.2024.0021>

Received: 5 January 2024

Accepted: 19 March 2024

Subject Category:

Ecology

Subject Areas:

ecology, behaviour, evolution

Keywords:

itinerant, life history, migration, reproduction,
shorebird

Author for correspondence:

Colby R. Slezak

e-mail: colby_slezak@uri.edu

Electronic supplementary material is available
online at <https://doi.org/10.6084/m9.figshare.c.7159242>.

Unconventional life history in a migratory shorebird: desegregating reproduction and migration

Colby R. Slezak¹, Erik J. Blomberg², Amber M. Roth², Liam A. Berigan², Alexander C. Fish², Rachel Darling², Sarah J. Clements², Greg Balkcom³, Bobbi Carpenter⁴, Gary Costanzo⁵, Jeffrey Duguay⁶, Clayton L. Graham¹, William Harvey⁷, Michael Hook⁸, Douglas L. Howell⁹, Seth Maddox¹⁰, Shawn W. Meyer¹¹, Theodore C. Nichols¹², J. Bruce Pollard¹³, Christian Roy¹⁴, Joshua C. Stiller¹⁵, Jacob N. Straub¹⁶, Mathieu Tetreault¹⁷, Reina Tyl¹⁸, Lisa Williams¹⁸, Jennifer E. Kilburn¹⁹ and Scott R. McWilliams¹

¹University of Rhode Island, Kingston, RI 02881, USA

²University of Maine, Orono, ME 04469, USA

³Georgia Department of Natural Resources, Wildlife Resources Division, Fort Valley, GA 31030, USA

⁴Florida Fish and Wildlife Conservation Commission, Gainesville, FL 32601, USA

⁵Virginia Department of Wildlife Resources, Charles City, VA 23188, USA

⁶Louisiana Department of Wildlife and Fisheries, Baton Rouge, LA 70898, USA

⁷Wildlife and Heritage Service, Department of Natural Resources, Cambridge, MD 21613, USA

⁸South Carolina Department of Natural Resources, Columbia, SC 29202, USA

⁹North Carolina Wildlife Resources Commission, Edenton, NC 27932, USA

¹⁰Alabama Department of Conservation and Natural Resources, Montgomery, AL 36130, USA

¹¹Environment and Climate Change Canada, Ottawa, Ontario, Canada K1V 1C7

¹²New Jersey Division of Fish and Wildlife, Woodbine, NJ 08270, USA

¹³Environment and Climate Change Canada, Sackville, New Brunswick, Canada E4L 1G6

¹⁴Environment and Climate Change Canada, Saint-Joseph Gatineau, Quebec, Canada K1A 0H3

¹⁵New York State Department of Environmental Conservation, Albany, NY 12233, USA

¹⁶State University of New York Brockport, Brockport, NY 14420, USA

¹⁷Environment and Climate Change Canada, D'Estimauville, Québec, QC, Canada G1J 0C3

¹⁸Pennsylvania Game Commission, Harrisburg, PA 17110, USA

¹⁹Rhode Island Department of Environmental Management, West Kingston, RI 02892, USA

CRS, 0009-0008-1266-4265; RD, 0000-0002-2457-8718; RT, 0000-0003-4540-778X; SRM, 0000-0002-9727-1151

Conventional life-history theory predicts that energy-demanding events such as reproduction and migration must be temporally segregated to avoid resource limitation. Here, we provide, to our knowledge, the first direct evidence of 'itinerant breeding' in a migratory bird, an incredibly rare breeding strategy (less than 0.1% of extant bird species) that involves the temporal overlap of migratory and reproductive periods of the annual cycle. Based on GPS-tracking of over 200 female American woodcock, most female woodcock (greater than 80%) nested more than once (some up to six times) with short re-nest intervals, and females moved northwards on average 800 km between first and second nests, and then smaller distances (*ca* 200+ km) between subsequent nesting attempts. Reliance on ephemeral habitat for breeding, ground-nesting and key aspects of life history that reduce both the costs of reproduction and migration probably explain the prevalence of this rare phenotype in woodcock and why itinerant breeding so rarely occurs in other bird species.

1. Introduction

Life-history theory predicts that resource limitations constrain when energy-demanding events such as reproduction occur and prohibits reproduction from overlapping with other energy-demanding events such as migration [1–3]. For most animals, timing of reproduction coincides with the period of the year when

Table 1. Avian species suspected of being itinerant breeders with the nature of the evidence (direct versus indirect) and associated citations.

common name	latin name	nature of the evidence ^a	citations
common quail	<i>Coturnix coturnix</i>	indirect	Moreau [26]
eared dove	<i>Zenaida auriculata</i>	indirect	Bucher [27]
Eurasian dotterel	<i>Charadrius morinellus</i>	indirect	Whitfield, in Wernham <i>et al.</i> [28]
yellow-billed cuckoo	<i>Coccyzus americanus</i>	indirect	Rohwer <i>et al.</i> [29]
hooded oriole	<i>Icterus cucullatus</i>	indirect	Rohwer <i>et al.</i> [29]
orchard oriole	<i>Icterus spurius</i>	indirect	Rohwer <i>et al.</i> [29], Rohwer <i>et al.</i> [30]
red-billed quelea	<i>Quelea quelea</i>	indirect	Ward [31], Jaeger <i>et al.</i> [32]
tri-coloured blackbird	<i>Agelaius tricolor</i>	indirect	Hamilton III [33], Wilson <i>et al.</i> [34]
phainopepla	<i>Phainopepla nitens</i>	indirect	Baldassarre <i>et al.</i> [35]
water rail	<i>Rallus aquaticus</i>	indirect	Lislevand <i>et al.</i> [36]
American woodcock	<i>Scolopax minor</i>	direct	Slezak <i>et al.</i> [37]

^a'Direct evidence' includes data on individuals that have resided in two distinct regions during the breeding season and were confirmed nesting in each region; 'indirect evidence' includes data on individuals that inhabit different regions during the breeding season but not confirmed breeding in both.

food resources are most available [4–6]. For species inhabiting seasonal environments, the timing of reproduction may have an outsized effect on breeding success and annual productivity [7–9], in part because food resources typically reach a seasonal peak in abundance and then decline over time [10–14]. Many migratory species time their northward migrations to coincide with food resources at important stopover locations to replenish fat reserves for migration and breeding [15]. Herbivores provide classic examples of such seasonal movements [16–18] including long-distance migratory geese that 'ride the green wave' north each spring as they prepare over time for breeding in the Arctic [19–21]. Across the diversity of life histories apparent in animals today, the vast majority of seasonal movements in migratory species conform to predictions of life-history theory in the sense that migration and reproduction occur during discrete periods and seldom overlap during the annual cycle in large part because of the substantial energy demands associated with breeding [3,22,23].

For migratory birds, balancing the timing of reproduction with that of migration has direct consequences: if individuals migrate too early in spring, their arrival may not coincide with peak resource abundance when raising young [6,24,25]; if they arrive too late, they may not have enough time to successfully raise young to independence prior to southward migration [25]. Only a few migratory bird species (less than 0.1% of extant species; table 1) have apparently defied conventional circannual patterns and overlap migratory and reproductive periods [23]. For these bird species, individuals breed then migrate and breed again, a breeding system known as 'itinerant breeding' or 'migratory double breeding' [29,35]. Careful consideration of such rare phenotypes may pose challenges to common assumptions about constraints on organic form and function [38] and allow us to identify ecological and physiological limitations that have produced them [39]. Although suspected in several species (table 1), evidence for this rare phenotype based on movement patterns of individuals has only been partly confirmed to date in three migratory bird species: redbilled quelea ([32] *Quelea quelea*), tri-coloured blackbird ([34] *Agelaius tricolor*) and phainopepla ([35] *Phainopepla nitens*). Direct evidence for itinerant breeding requires tracking individual movements during the breeding season while confirming breeding in multiple regions; such evidence is not available to date for any of these purported itinerant breeders. Herein, we provide, to our knowledge, the first direct evidence of itinerant breeding from detailed GPS tracking of greater than 200 individually marked female American woodcock (*Scolopax minor*), a migratory shorebird that inhabits forests in eastern and central North America.

We closely monitored the movement patterns of GPS-tagged female woodcock across their entire range over two years (2021–2022) to document migration movements as well as nest initiation and incubation. For 26 females that exhibited short (e.g. <20 m) step-lengths in sequence over several days (electronic supplementary material, figure S1), we field-verified that these distinct movement patterns were associated with an actively nesting individual. We then used the nestR package [40] to retroactively locate nest attempt(s) for a larger sample of females ($n = 272$) that were captured and tagged over 4 years (2019–2022); we were also able to confirm correct classification of 92% (24 of 26) of our field-verified nesting attempts when we used this approach (see Methods). A subset of females ($n = 55$) were tracked frequently (GPS locations every 1–3 days; $\bar{x} = 4.3$ locations every 4 days) over an extended portion of the breeding season (20 February to 1 June). For these individuals we were able to evaluate the overlap between migration and reproduction, estimate key components of their reproductive strategy (i.e. number of nest attempts, attempt duration and re-nest time), and determine how this strategy differed for young versus older woodcock (*ca* 1 yr old versus ≥ 2 yr old females). If woodcock are itinerant breeders, then we predicted that females would regularly and rapidly re-nest after nest failure, that females would migrate north between nesting attempts, and in general there would be considerable temporal overlap between the reproduction and migration periods of the annual cycle.

2. Methods

(a) Satellite transmitter attachment

Thirty-six state, federal and non-governmental organizations helped tag female woodcock throughout eastern North America from Florida, USA northwards to Atlantic Canada (electronic supplementary material, table S1) as part of the Eastern Woodcock Migration

Research Cooperative [41]. Woodcock were captured and leg-banded at night using the night lighting technique [42] or mist-netted during crepuscular periods [43]. We determined age and sex of woodcock in the field using wing feather characteristics [44]. From 1 September 2019 to 1 June 2022, collaborators captured 366 females that were then tagged with 4 g, 5 g or 6 g GPS satellite transmitters (PinPoint Argos 100; PinPoint Argos 125; PinPoint Argos 150; Lotek Wireless, Newmarket, Ontario, Canada) that collected 75 to 150 GPS locations before tag failure. We closely followed protocols outlined in the *Guidelines to the use of Wild Birds in Research* to minimize stress and disturbance of captured birds and we ensured that total mass of marking materials (transmitter, harness and band) did not exceed 4% body mass [45]. The 272 tags still collecting data on or after 1 January annually (electronic supplementary material, table S1) were programmed to transmit GPS locations (± 20 m) approximately every 1–3 days, although the actual number of locations recorded was on average 4.3 locations (s.d. = 5.42) every 5 days for the period of time tags were active. The collected locations from all PinPoint tags were uploaded to our Movebank repository.

(b) Field verification: sequential movement distances indicate nesting females

During 2021–2022, we scrutinized the daily GPS locations of females during the window of time encompassing the bulk of courtship activity from 20 February to 1 June each year. Specifically, we measured the step-lengths between subsequent GPS locations for each female in the dataset at least twice a week using the ‘as.traj’ function in the *adehabitatLT* package [46] in R version 4.2.2 [47]. We recognize that some southerly nesting woodcock may initiate nests earlier than mid-February; however, many tags were still being deployed in the southerly areas during early-February and thus did not start transmitting until mid- to late- February (electronic supplementary material, table S1). When step-lengths were small (less than 20 m) for two to three or more GPS transmissions (electronic supplementary material, figure S1), we notified the nearest collaborator and requested they travel to the point cluster and investigate the site for a nesting female or signs of a successful or failed nesting attempt (e.g. longitudinally split eggshells or crushed eggshells). In total, we field-verified 26 nests (2021: $n = 16$; 2022: $n = 10$) from 22 (2021: $n = 13$; 2022: $n = 9$) individual females including four females confirmed nesting more than once. The step lengths for field-verified nesting females in 2021 (12.6 m, 95% confidence interval (CI): 11.9–13.3 m) were not statistically different ($F_{1,178} = 0.43$, $p = 0.52$) from 2022 (14.2 m, 95% CI: 12.8–15.6 m) and confirmed that these short step-lengths between consecutive *ca* daily locations indicated an incubating female woodcock. In addition, there were 11 instances that met these step-length criteria for incubating females but cooperators were unable to find the nest. Such potential false-positive detections probably occurred because female woodcock and their nests are highly cryptic and so easily missed or because nest depredation rates can be high for ground-nesting woodcock ([48] reports 67% failure) and so nests may have gone before cooperators were asked to find the nest. Three of the four females that were found re-nesting migrated a substantial distance northwards from their initial nest attempt (range: 199 to 889 km).

(c) Application to larger (2019–2022) GPS tracking dataset

Given that our tracking data covered the majority of breeding and wintering range of woodcock in North America, we used a more inclusive (1 January to 1 September) as well as restrictive (20 February to 1 June) time period to investigate nesting behaviour of females. Our more inclusive dataset allowed us to include more females that were tracked for variable durations of time and so evaluate population-level patterns of nesting in relation to timing and latitude. By contrast, our more restrictive dataset included only females that were tracked almost daily (on average 3.5 locations every 4 days) from 20 February to 1 June, which includes the approximate range of core breeding activity in the most southerly and northern reaches of the woodcock’s range [43,49–51]. These females tracked throughout the core breeding period allowed us to answer fundamental questions about woodcock breeding ecology such as the total number of nesting attempts, how long nest attempts last on average, how quickly females re-nest, if distance moved between attempts was related to female age, and whether migration distance influenced total number of nest attempts. This dataset also allowed us to relate a female’s nesting sequence to metrics such as distance moved, latitude and week of nesting season.

(d) Population-level patterns of nesting throughout the breeding season (1 January to 1 September)

We used the `find_nests()` function in the *nestR* package [40] to identify nesting attempts from our inclusive and restrictive datasets, and used our sample of field-verified nests to assess its accuracy. The `find_nests()` function identifies potential nests based on revisitations and the amount of time spent within a user specified buffer surrounding each GPS location [40]. Our inclusive dataset included a broader range of start (1 January) and end (1 September) dates for the four nesting seasons (2019–2022) to ensure early- or late-nesting attempts by females were detected [52–54]. Because the step lengths of our field-verified nesting females were on average < 20 m in each year, we chose a maximum buffer size of 20 m that females had to stay within for at least 5 days to be considered a nesting attempt. We chose a 5-day window because the satellite tags used in this project require about 5 days to collect three successful GPS locations and transmit to the ARGOS satellite, thus this was typically the minimum number of days over which nesting females could be detected. Woodcock take infrequent incubation recesses [55] and our transmitter schedule was coarse enough that capturing a female off of the nest was unlikely, thus we set the ‘`min_d_fix`’ parameter to two GPS points, which required two consecutive GPS fixes to fall outside of a 20 m buffer before a nest was considered no longer active. This allowed us to approximate nest attempt start and end dates for each of the female’s nesting attempts. When using the ‘`find_nests`’ function, we set the ‘`discard_overlapping`’ parameter to ‘TRUE’ to ensure that only one location would be chosen from a set of temporally overlapping nest attempts. For females that nested more than once, we measured the distance travelled to subsequent nesting attempts using the ‘`as.traj`’ function in the *adehabitatLT* package [46] in R.

Our use of short-distance step lengths to identify incubating females could unintentionally include stationary females that had died, and so the following two-step process was used to exclude ‘nests’ from females after they were classified as deceased. We first used the *momentuHMM* package in R [56] to train a hidden Markov model on a set of known movement states (dead female, alive female but stationary and migratory females) and used this trained model to autonomously classify mortality events in our female tracking dataset. Once the model had identified which females had probably died, we used a R Shiny application to ensure that any individual marked as deceased had (i) 15 locations in a deceased state, and (ii) at least half of deceased locations fell within a threshold distance of the centroid. The threshold distance varied based on landcover owing to differences in GPS accuracy between cover types (range of threshold distances: 3.93–10.85 m). This two-step process identified and excluded 13 females that would have been classified as ‘nesting’ but instead had died during the four breeding seasons.

(e) Individual-level patterns of nesting during the core breeding period (20 February to 1 June)

Of the 272 females tracked from 1 January to 1 September, a subset ($n=55$ females) with greater than 10 GPS locations were tracked throughout the entire period of core breeding activity observed in the southern and northern reaches of the woodcock's breeding range (20 February to 1 June; [52–54,57]) during at least one of the 4 years of the study. We used this subset of 55 females to evaluate the total number of nesting attempts by each female in a given season, the date and estimated duration of each nesting attempt, and how quickly females re-nested. Like the inclusive dataset, we measured the distance moved for females that nested more than once using the 'as.traj' function in the adehabitatLT package [46] in R, and then related distance moved between attempts to female age. This dataset also allowed us to relate a female's nesting sequence to metrics such as distance moved, latitude and week of nesting season because females had complete tracking data (i.e. the entirety of the core breeding period) unlike the females in the more inclusive dataset that were often tracked for a more brief and variable period of time.

(f) Statistical methods

(i) Population-level patterns of nesting throughout the breeding season (1 January to 1 September)

We calculated nest attempt duration for all nests by subtracting the estimated nest attempt end date by the nest attempt start date using the 'difftime' function in R. Nests that lasted 19 or more days were considered likely to be successful (mean incubation period is 21 days for precocial woodcock; [58], range is 19 to 22 days; [49]). To determine if week of nesting season, latitude, or the additive effect of these two variables best predicted nest success, we used logistic regression models implemented using a generalized linear model and logit-link function in the stats package in R. However, we did not include the additive model in our final model comparison because latitude was not a significant predictor. We performed model selection using Akaike information criterion adjusted for small sample sizes (AIC_c) using the 'aictab' function in the AICcmodavg package [59]. We considered models with ΔAIC_c values ≤ 2 to have equivalent evidence of being the top model(s) [60]. Finally, we used the 'geom_smooth' function in the ggplot2 package in R to plot a smoothed trend line (loess line) so that we could visualize the relationship of ordinal date and latitude with the distance moved between nesting attempts for our entire dataset of tagged females.

(ii) Individual-level patterns of nesting during the core breeding period (20 February to 1 June)

For each of the 55 females tracked throughout the core breeding period, we also calculated nest attempt duration for every documented nest by subtracting nest attempt end date by nest attempt start date using the 'difftime' function in R. We also used the 'difftime' function to calculate re-nest time for females that nested more than once by subtracting the attempt start date by the attempt end date of the prior nesting attempt. We conducted an analysis of variance (ANOVA) using the 'aov' function in R to determine whether nest attempt duration and re-nest time differed among the first, second, third or fourth nest attempt. We had too few fifth ($n=6$) and sixth ($n=1$) nesting attempts to include them for this statistical analysis. We also used an ANOVA to determine if distance travelled to subsequent nest attempt varied by female age (second year (SY) versus after second year (ASY)) or nest attempt sequence (1–4).

3. Results

(a) Population-level patterns: nesting occurs from January–August, and females move north between nesting attempts

We identified 336 woodcock nesting attempts from 153 of 272 (56.25%) females that transmitted locations between 1 January and 1 September during the 4 years (i.e. 2019 to 2022). The 119 females that were not detected nesting were typically tracked for short periods of time on average (43.39, 95% CI: 37.34–49.43 days) and half of these 119 females' tags had 35 or less transmissions, a period too short to reliably detect if these females attempted nesting. Like Whiting [54], we found that a majority of nests were initiated no earlier than mid-February; only 72 (22%) of 336 were initiated prior to 20 February, all in more southerly areas of the woodcock's range ([61] reports Alabama, Arkansas, Florida, Georgia, Louisiana, Maryland, North Carolina, New Jersey, Oklahoma, South Carolina and Virginia). Females that nested earlier in the spring tended to move further to subsequent nesting attempts than those later in the season (figure 1a). In addition, females tended to move a shorter distance (km) between subsequent nesting attempts as latitude increased (figure 1b). We found that about half (57%) of all nesting attempts were long enough in duration to have hatched chicks (≥ 19 days incubation) although such an estimate of nest survival is inflated because our methodology limited our ability to detect short-lived nests (i.e. lasting ≤ 3 days or before the tag collected three GPS transmissions). An incubation period of 19 or more days also does not necessarily equate to successfully fledged chicks because brood loss is not considered. Our top model ($\Delta AIC_c \leq 2$) with week of nesting season predicting nest success held 86% of the cumulative weight of our three model sets (electronic supplementary material, table S2). This model suggests that the probability of a successful nesting attempt increased (week coefficient = 0.05, 95% CI: 0.018–0.083) as the nesting season progressed.

(b) Individual-level patterns: females first nest on 'wintering areas' and often re-nest many times (up to 6) as they migrate north

We identified 144 likely nesting attempts from 52 of the 55 females (95%) that were tracked every 1 to 3 days throughout the core breeding season (20 February to 1 June in a given year), and 42 of these 52 females (81%) had more than one nesting attempt (figure 2a). We found no statistical difference ($F_{3,133}=1.86$, $p=0.14$) in nest duration between nesting attempts 1 to 4 (figure 2b), so we report the population mean (16.47, 95% CI: 15.65–183 days). We also found no statistical difference ($F_{2,82}=2.46$, $p=0.092$) in re-nest time between nesting attempts 1 to 4 (figure 2c) so we report the population mean (23.95, 95% CI: 20.04–27.86 days; range: 1–70 days). However, distance moved between nesting attempts 1 to 4 (figure 2d) significantly differed




Figure 1. Movement patterns between nesting attempts for a population of individually marked females. (a) Smoothed loess line with 95% confidence bands depicting the moving average of distance moved between nesting attempts by week of nesting season for 337 nesting attempts from 154 females tracked annually from 2019 to 2022. (b) Smoothed loess line with 95% confidence bands depicting the moving average of distance moved between nesting attempts by latitude for 337 nesting attempts from 154 females tracked annually from 2019 to 2022.




Figure 2. Nesting characteristics for individually marked females. (a) Total number of females with one or more nest attempts, (b) nest attempt duration, (c) re-nesting interval time between each nest attempt, and (d) distance moved between nesting attempts for 144 nesting attempts for 52 of 55 female American woodcock that nested and were tracked almost daily from 20 February to 1 June in a given year (2019–2022). Boxes indicate median, mean (diamond in grey), 25% and 75% quartiles. Points represent raw data values, and whiskers represent the greatest and least values within 1.5 times the interquartile range over the 75th percentile. Outliers are presented as a cross-hatch (X).

($F_{2,81} = 10.42$, $p < 0.05$). Females moved further between nesting attempt 1 to 2 than their subsequent nesting attempts regardless of location or latitude although the mean distance travelled after attempts 2 and 3 did not statistically differ (mean of attempt 1 to 2 (799.05 km, 95% CI: 620.81–977.29 km); mean of attempt 2–3 (330.62 km, 95% CI: 128.37–532.88.17 km); mean of attempt 3–4




Figure 3. Nest sequence by week and latitude for individually marked females. (a) Raincloud plot showing the distribution of nest initiation dates by week and (b) by latitude for each subsequent nesting attempt (up to five attempts) for the 144 nesting attempts for 52 of 55 female American woodcock that nested and were tracked almost daily from 20 February to 1 June in a given year (2019–2022). Boxes indicate median, mean (black point within box plot), 25% and 75% quartiles. Points represent raw data values, and whiskers represent the highest and lowest values within 1.5 times the interquartile range over the 75th percentile.

(205.74, 95% CI: 10.89–400.60 km)). Distance moved between subsequent nesting attempts did not statistically differ for SY versus ASY females ($F_{1,77} = 3.31$, $p = 0.07$) although older ($n = 44_{ASY}$) females moved further on average (619.20 km, 95% CI: 432.48–805.91 km) between attempts than younger ($n = 71_{SY}$) females (388.84 km, 95% CI: 270.42–507.25 km). Most first nests occurred before 12 March (figure 3a) and were below 40° N latitude (range: 30° N to 48° N latitude; figure 3b) within areas outside of the area traditionally reported as the woodcock breeding range [58]. We found that woodcock had a prolonged nesting season, with some nests initiated as early as 9 January and as late as 10 August, and that females nesting at higher latitudes were typically those that had already nested at lower latitudes (figure 3b). Between 1 January and the final tag transmission, each female migrated northwards on average 1348.28 km (95% CI: 1210.68–1485.87 km); however, the total distance a female migrated did not relate to its total number of nesting attempts ($F_{1,53} = 0.86$, $p = 0.36$). Many females first nested in the southern portion of the woodcock's range (mean 38.03° latitude, 95% CI: 36.85–39.21° latitude), and then continued northwards to nest again in a direction typical of migratory woodcock (electronic supplementary material, figure S2).

4. Discussion

As we have shown for American woodcock, itinerant breeders can rapidly switch between reproductive and migratory states and this challenges the fundamental assumption that endocrine mismatch forces divergent partitioning of certain life-history traits (i.e. reproduction and migration) across the annual cycle [30,35]. Flexible timing in the annual cycle of birds is thought to arise when the breeding season is prolonged (e.g. non-migrants that are not constrained by periods of migration) and when there is great annual variation in environmental cues (e.g. climatic conditions and shifting food availability) which makes consistent timing of reproduction less beneficial [35,62]. However, we do not think this set of factors alone adequately explains the unusual breeding strategy of woodcock and potentially the other purported itinerant breeders that have unconstrained themselves from distinct breeding sites and seem to migrate between breeding attempts (table 1).

Itinerant breeding birds documented to date share the following traits: a relatively low cost of reproduction and reliance on ephemeral habitats (e.g. early successional forest for American woodcock [23,35]). These traits are important in explaining itinerant breeding in woodcock, but other well understood aspects of the woodcock's life history seem potentially relevant in explaining how this could evolve in woodcock and potentially other itinerant breeders (figure 4). Like many shorebirds, woodcock consistently lay four eggs and produce precocial young at hatch [63]. Costs of reproduction and migration for female woodcock are estimated to be relatively small compared to other Arctic breeding shorebirds, and like other shorebirds females typically rely on exogenous nutrients acquired upon arrival to breeding sites for egg production [64,65]. However, woodcock are relatively unusual shorebirds in that females also store appreciable fat in preparation for reproduction [64,66]. The considerably shorter distance migrations than that of most other shorebirds along with their reliance on exogenous nutrients may explain why females can re-nest frequently even while providing uniparental care [67]. Woodcock are among the earliest breeding




Figure 4. Life-history traits and ecological conditions related to the occurrence of a rare breeding strategy, itinerant breeding, in migratory birds. Itinerant breeding involves desegregating in time the reproduction and migration periods such that nesting occurs multiple times while migrating between nest attempts. Selection for itinerant breeding among migratory birds requires reduced costs of migration and reproduction along with a mating system that permits mate finding along migratory routes and the ability to re-nest. For woodcock, females alone build the ground-nest, lay the initial four-egg clutch, incubate and tend the precocial young. Reversed sexual size dimorphism is pronounced in woodcock and thus for the large-bodied females the costs of producing a clutch is relatively low. Like many ground-nesting birds, nest predation is common, and thus the ability to quickly re-nest and prolong the breeding season is a key determinant of lifetime reproductive success. What makes itinerant breeders quite unusual is that instead of choosing to re-nest nearby, they migrate between nesting attempts. Woodcock initiate breeding very early each year (January/February) often at or near wintering areas with males courting and seeking mating opportunities throughout their migration to northern breeding areas. Their polygynous mating system along with the relatively low costs of reproduction allow female woodcock to find mates and re-nest several times (up to six) as they track spring resource pulses northwards during spring migration.

birds in the Northern Hemisphere and commence nesting when environmental conditions are often not favourable for other species [68,69]. As a consequence, woodcock are more susceptible to unpredictable environmental conditions that could lead to nest failure (i.e. late frosts, spring snow/ice storms and summer heat waves) than species with shorter reproductive windows. This phenological pattern is similar to the phainopepla system where frost or drought can kill the mistletoe crop and disrupt early breeding [35]. An extended reproductive period, with the opportunity for multiple re-nesting attempts may allow woodcock to better cope with nest failure associated with stochastic weather events as well as predation. The unusually large size of female woodcock relative to males [70] and a total fresh egg mass that is a relatively small percentage of the female's mass [71], combined with a simple nest (scrape lined with leaf litter) and precocial young [58] reduces the relative cost of each nesting attempt and may facilitate the rapid (figure 2c), frequent (figure 2a), and geographically expansive re-nesting behaviour (figure 2d) of this itinerant breeder. The reduced costs of each nesting attempt combined with their reliance on exogenous nutrients for egg laying may also select for woodcock to track seasonally abundant food and nesting cover as they 'follow the spring thaw north' (figure 5) much like avian herbivores that 'ride the green wave' as they accumulate fat stores on their way to high-latitude breeding areas [16,17].

The recognition that woodcock have evolved an itinerant breeding strategy also helps explain some of the long-recognized, unusual aspects of the life history of this species: (i) extended breeding seasons (i.e. nesting January to August [52,54,72,73]) with males initiating courtship displays as early as January in the south [51,74], (ii) re-nesting as a rule rather than an exception [67] and its frequency declining in years with late spring green-up [75], (iii) females that exhibited peculiar movement patterns (e.g. brooding females banded in the south later harvested in the far north of their range [76]), and (iv) extreme levels of reverse sexual size dimorphism compared to other shorebirds [70]. The breeding system of woodcock has been characterized as polygynous with males defending territories in which they perform courtship displays and females nesting near or in these territories, but with no formation of a social pair bond, and no evidence of multiple paternity within broods [58]. Unlike prior studies, we directly tracked individual females to reveal multiple nesting attempts interrupted by long-distance movements along their northly migratory paths in spring (figure 5). This has allowed us to confirm itinerant breeding at the individual (figure 2) and population level




Figure 5. Example movement patterns for individually marked nesting females during spring migration. The location of nests for the few females that nested only once ($n = 10$) are displayed as black diamonds. Lines connecting nesting locations (circles) provide an example GPS track for a female that re-nested twice (yellow), three times (green), four times (blue), five times (red) and six times (purple). Arrows show the northerly direction of female movement between each nest attempt. Arrowheads without a visible line segment were re-nests that were short distances from the previous nest attempt (typically ≤ 5 km), and the direction of the arrowhead indicates the direction moved from the previous nesting attempt. The black dashed line at 40° N latitude represents the southern extent of the area considered the core region of woodcock breeding activity. GPS tracks of all females that nested ($n = 52$) are provided in the electronic supplementary material, figure S2.

(figure 1). Prior studies of itinerant breeders (table 1) have been unable to identify the locations of multiple breeding attempts and have instead relied on the movement patterns of a few individuals which had presumably nested before migrating to a new breeding site [35]. The large proportion of woodcock that re-nested (figure 2a; 81% of all nesting females) and their impressive migratory movements (up to 2315 km) between breeding attempts (electronic supplementary material, figure S2), combined with the movement patterns and nesting attempts observed in the entire population (figure 1), suggest that itinerant breeding is prolific among woodcock. For woodcock, we hypothesize that certain aspects of their life history (i.e. reverse sexual size dimorphism, clutch size limitations, precocial young, mixed capital-income breeding strategy [6] have reduced overall reproductive cost, allowing them to nest in early-successional forests over extended periods of time and track thawing soils and seasonally abundant nesting cover during spring. It remains to be determined if these ecological and life-history factors also explain itinerant breeding in other species and how migratory species at large adapt to changing environments.

Ethics. All woodcock trapping, handling and tagging activities were conducted in accordance with the University of Rhode Island Animal Care and Use Guidelines under Animal Care and Use Protocol AN10A02A017 and the University of Maine Institutional Animal Care and Use Committee (protocols A2017_05_02 and A2020_07_01).

Data accessibility. All female capture information, GPS tracking data downloaded from Movebank and R code is available online in our Zenodo repository [37]. Supplementary material is available online [77].

Declaration of AI use. We have not used AI-assisted technologies in creating this article.

Authors' contributions. C.R.S.: conceptualization, data curation, formal analysis, investigation, methodology, visualization, writing—original draft, writing—review and editing; E.J.B.: conceptualization, data curation, funding acquisition, investigation, methodology, resources, supervision, writing—review and editing; A.M.R.: funding acquisition, writing—review and editing; L.A.B.: investigation, visualization, writing—review and editing; R.D.: investigation, visualization, writing—review and editing; A.C.F.: conceptualization, investigation, writing—review and editing; S.J.C.: writing—review and editing; G.B.: investigation; B.C.: investigation; G.C.: investigation; J.D.: investigation; C.L.G.: investigation; W.H.: investigation; M.H.: investigation; D.L.H.: investigation; S.M.: investigation; S.W.M.: investigation; T.C.N.: investigation; J.B.P.: investigation; C.R.: investigation, writing—review and editing; J.C.S.: investigation; J.N.S.: investigation, writing—review and editing; R.T.: writing—review and editing; L.W.: investigation; J.E.K.: investigation; M.T.: investigation, writing—review and editing; S.R.M.: conceptualization, funding acquisition, investigation, methodology, supervision, visualization, writing—original draft, writing—review and editing.

All authors gave final approval for publication and agreed to be held accountable for the work performed therein.

Conflict of interest declaration. We declare we have no competing interests.

Funding. Funding was provided from the U.S. Fish and Wildlife Service Webless Migratory Game Bird Research Fund (grant no. F), the US Department of Agriculture McIntire-Stennis grant nos ME0-21422, ME0-q 42108 and MS-983, the US Department of Agriculture Hatch (grant no. H-338) and the Rhode Island Department of Environmental Management via U.S. Fish and Wildlife Service and Sport Fish Restoration Program (grant no. W-23R).

Acknowledgements. We thank the collective group of talented field biologists, researchers and volunteers who made field-verification of nests possible: J. Carloni, T. Cunningham, S. Lamonde, G. Champagne, P. Gilboe, D. Sausville, L. Simulik, R. Feldman, L. Millett, A. Hanson, W. Wolanske, J. McDonald, T. Young, C. Baranski, H. Wells, T. Buchanan, K. Connor and S. Heerkens.

References

1. Roff DA, Fairbairn DJ. 2007 The evolution of trade-offs: where are we? *J. Evol. Biol.* **20**, 433–447. (doi:10.1111/j.1420-9101.2006.01255.x)
2. Roff DA, Heibo E, Vøllestad LA. 2006 The importance of growth and mortality costs in the evolution of the optimal life history. *J. Evol. Biol.* **19**, 1920–1930. (doi:10.1111/j.1420-9101.2006.01155.x)
3. Stearns SC. 1992 *The evolution of life histories*. Oxford, UK: Oxford University Press.
4. Gustine D, Barboza P, Adams L, Griffith B, Cameron R, Whitten K. 2017 Advancing the match-mismatch framework for large herbivores in the Arctic: evaluating the evidence for a trophic mismatch in caribou. *PLoS ONE* **12**, e0171807. (doi:10.1371/journal.pone.0171807)
5. Evans SR, Bearhop S. 2022 Variation in movement strategies: capital versus income migration. *J. Anim. Ecol.* **91**, 1961–1974. (doi:10.1111/1365-2656.13800)
6. Drent RH, Daan S. 1980 The prudent parent: energetic adjustments in avian breeding. *Ardea* **55**, 225–252.
7. Post E, Forchhammer MC, Stenseth NC, Callaghan TV. 2001 The timing of life-history events in a changing climate. *Proc. R. Soc. B* **268**, 15–23. (doi:10.1098/rspb.2000.1324)
8. Réale D, McAdam AG, Boutin S, Berteaux D. 2003 Genetic and plastic responses of a northern mammal to climate change. *Proc. R. Soc. B* **270**, 591–596. (doi:10.1098/rspb.2002.2224)
9. Durant JM, Hjermmann DØ, Ottersen G, Stenseth NChr. 2007 Climate and the match or mismatch between predator requirements and resource availability. *Clim. Res.* **33**, 271–283. (doi:10.3354/cr033271)
10. Verboven N, Visser ME. 1998 Seasonal variation in local recruitment of great tits: the importance of being early. *Oikos* **81**, 511–524. (doi:10.2307/3546771)
11. Naef-Daenzer B, Widmer F, Nuber M. 2001 Differential post-fledging survival of great and coal tits in relation to their condition and fledging date. *J. Anim. Ecol.* **70**, 730–738. (doi:10.1046/j.0021-8790.2001.00533.x)
12. Verhulst S, Nilsson J-Å. 2008 The timing of birds' breeding seasons: a review of experiments that manipulated timing of breeding. *Phil. Trans. R. Soc. B* **363**, 399–410. (doi:10.1098/rstb.2007.2146)
13. Gruebler MU. 2010 Fitness consequences of timing of breeding in birds: date effects in the course of a reproductive episode. *J. Avian Biol.* **41**, 282–291. (doi:10.1111/j.1600-048X.2009.04865.x)
14. Perrins CM. 1965 Population fluctuations and clutch-size in the great tit, *Parus major* L. *J. Anim. Ecol.* **34**, 601–647. (doi:10.2307/2453)
15. Dingle H. 2014 Seasonal migration patterns. In *Migration: the biology of life on the move* (ed. H Dingle), pp. 163–182. Oxford, UK: Oxford University Press.
16. Middleton AD, Merkle JA, McWhirter DE, Cook JG, Cook RC, White PJ, Kauffman MJ. 2018 Green-wave surfing increases fat gain in a migratory ungulate. *Oikos* **127**, 1060–1068. (doi:10.1111/oik.05227)
17. Merkle JA *et al.* 2016 Large herbivores surf waves of green-up during spring. *Proc. R. Soc. B* **283**, 20160456. (doi:10.1098/rspb.2016.0456)
18. Abraham JO, Upham NS, Damian-Serrano A, Jesmer BR. 2022 Evolutionary causes and consequences of ungulate migration. *Nat. Ecol. Evol.* **6**, 998–1006. (doi:10.1038/s41559-022-01749-4)
19. Drent R, Ebbinge B, Weijand B. 1979 Balancing the energy budgets of Arctic-breeding geese through-out the annual cycle: a progress report. *Verh. Orn. Ges. Bayern* **23**, 239–264.
20. Shariatinajafabadi M *et al.* 2014 Migratory herbivorous waterfowl track satellite-derived green wave index. *PLoS ONE* **9**, e10831. (doi:10.1371/journal.pone.0108331)
21. Si Y, Xin Q, De Boer WF, Gong P, Ydenberg RC, Prins HHT. 2015 Do Arctic breeding geese track or overtake a green wave during spring migration? *Sci. Rep.* **5**, 8749. (doi:10.1038/srep08749)
22. Stearns SC. 1989 Trade-offs in life-history evolution. *Funct. Ecol.* **3**, 259–268. (doi:10.2307/2389364)
23. Newton I. 2008 *The migration ecology of birds*, 1st edn. Oxford, UK: Academic Press.
24. Salido L, Purse BV, Marrs R, Chamberlain DE, Shultz S. 2012 Flexibility in phenology and habitat use act as buffers to long-term population declines in UK passerines. *Ecography* **35**, 604–613. (doi:10.1111/j.1600-0587.2011.06797.x)
25. Åkesson S, Ilieva M, Karagicheva J, Rakhimberdiev E, Tomotani B, Helm B. 2017 Timing avian long-distance migration: from internal clock mechanisms to global flights. *Phil. Trans. R. Soc. B* **372**, 20160252. (doi:10.1098/rstb.2016.0252)
26. Moreau RE 1951 The British status of the quail and some problems of its biology. *Brit. Birds* **44**, 257–276.
27. Bucher EH 1982 Colonial breeding of the eared dove (*Zenaidura macroura*) in northeastern Brazil. *Biotropica* **14**, 255.
28. Wernham CV, Toms MP, Marchant JH, Clark JA, Siriwardena GM, Baillie SR. (eds) 2002 *The migration atlas: movements of the birds of Britain and Ireland*. London, UK: T. & A.D. Poyser.
29. Rohwer S, Rohwer VG, Peterson AT, Navarro-Sigüenza AG, English P. 2012 Assessing migratory double breeding through complementary specimen densities and breeding records. *Condor* **114**, 1–14. (doi:10.1525/cond.2012.110064)
30. Rohwer S, Hobson KA, Rohwer VG. 2009 Migratory double breeding in neotropical migrant birds. *Proc. Natl Acad. Sci. USA* **106**, 19 050–19 055. (doi:10.1073/pnas.0908121106)
31. Ward P. 1971 The migration patterns of *Quela quelea* in Africa. *Ibis* **113**, 275–297.
32. Jaeger MM, Bruggers RL, Johns BE, Erickson WA. 1986 Evidence of itinerant breeding of the red-billed quelea quelea quelea in the Ethiopian Rift Valley. *Ibis* **128**, 469–482. (doi:10.1111/j.1474-919X.1986.tb02699.x)
33. Hamilton III WJ. 1998 Tricolored blackbird itinerant breeding in California. *Condor* **100**, 218–226.
34. Wilson CR, Meese RJ, Wyckoff AC. 2016 Breeding chronology, movements, and life history observations of tricolored blackbirds in the California Central Coast. *Calif. Fish Game* **102**, 162–174.
35. Baldassarre DT, Campagna L, Thomassen HA, Atwell JW, Chu M, Crampton LH, Fleischer RC, Riehl C. 2019 GPS tracking and population genomics suggest itinerant breeding across drastically different habitats in the phainopepla. *Auk* **136**, 1–12. (doi:10.1093/auk/ukz058)
36. Lislevand T, Hahn S, Rislää S, Briedis M. 2020 First records of complete annual cycles in water rail *Rallus aquaticus* show evidence of itinerant breeding and a complex migration system. *J. Avian Biol.* **51**, 1–8.
37. Slezak CR. 2024 Code for: Unconventional life-history in a migratory shorebird: desegregating reproduction and migration. *Zenodo*. (doi:10.5281/zenodo.10037878)
38. Dudley R, Vermeij GJ. 1992 Do the power requirements of flapping flight constrain folivory in flying animals? *Funct. Ecol.* **6**, 101–104.
39. Vermeij GJ. 1987 *Evolution and escalation: an ecological history of life*. Princeton, NJ: Princeton University Press.
40. Picardi S *et al.* 2020 Analysis of movement recursions to detect reproductive events and estimate their fate in central place foragers. *Mov. Ecol.* **8**, 1–14. (doi:10.1186/s40462-020-00201-1)
41. Blomberg E, Roth A, Fish A. 2023 Eastern Woodcock Migration Research Cooperative. See woodcockmigration.org (accessed 8 March 2023).
42. Shuler JF, Samuel DE, Shissler BP, Ellingwood MR. 1986 A modified nightlighting technique for male American woodcock. *J. Wildl. Manag.* **50**, 384–387. (doi:10.2307/3801091)

43. McAuley DG, Longcore JR, Sepik GF. 1993 Techniques for research into woodcocks: experiences and recommendations. In *Proc. of The Eighth American Woodcock Symp.* (eds JR Longcore, GF Sepik), pp. 5–11. Washington, DC: U.S. Fish and Wildlife Service.
44. Martin FW. 1964 Woodcock age and sex determination from wings. *J. Wildl. Manag.* **28**, 287–293. (doi:10.2307/3798090)
45. Fair J, Paul E, Bies L (eds) 2023 *Guidelines to the use of wild birds in research*. See <http://www.BIRDNET.org>.
46. Calenge C. 2006 The package 'adehabitat' for the R software: a tool for the analysis of space and habitat use by animals. *Ecol. Modell.* **197**, 516–519. (doi:10.1016/j.ecolmodel.2006.03.017)
47. R Core Team. 2022 *R: a language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing.
48. Huinker AE. 2020 American woodcock habitat selection and reproductive success in Michigan. PhD thesis, Michigan State University, East Lansing, MI, USA.
49. Sepik GF, Owen RB, Coulter MW. 1981 A landowner's guide to woodcock management in the northeast. In *U.S. Fish and Wildlife Service Miscellaneous Report 253*, Calais, ME: U.S. Fish and Wildlife Service.
50. Murphy DW, Thompson FR. 1993 Breeding chronology and habitat of the American woodcock in Missouri. In *Proc. of The Eighth American Woodcock Symp.* (eds JR Longcore, GF Sepik), pp. 12–18. Washington, DC: U.S. Fish and Wildlife Service.
51. Tappe PA, Whiting RM, George RR. 1989 Singing-ground surveys for woodcock in east Texas. *Wildl. Soc. Bull.* **17**, 36–40.
52. Hicks LE. 1933 The breeding birds of Ashtabula County. *Wilson Bull.* **45**, 168–195.
53. Pitelka FA. 1943 Territoriality, display, and certain ecological relations of the American woodcock. *Wilson Bull.* **55**, 88–114.
54. Whiting MR, Haukos DA, Smith LM. 2005 Factors affecting January reproduction of American woodcock in Texas. *Southeastern Nat.* **4**, 639–646. (doi:10.1656/1528-7092(2005)004[0639:FAJROA]2.0.CO;2)
55. McAuley DG, Glugston DA, Halteman WA. 2019 Behavior of incubating American woodcock (*Scolopax minor*) in Maine. In *Proc. of the Eleventh American Woodcock Symp.* (eds DG Krementz, DE Andersen, TR Cooper), pp. 265–269. Roscommon, MI: University of Minnesota Libraries Publishing.
56. McClintock BT, Michelot T. 2018 momentuHMM: R package for generalized hidden Markov models of animal movement. *Methods Ecol. Evol.* **9**, 1518–1530. (doi:10.1111/2041-210X.12995)
57. Roberts TH. 1993 The ecology and management of wintering woodcocks. In *Proc. of the Eighth American Woodcock Symp.* (eds JR Longcore, GF Sepik), pp. 87–97. Washington, DC: U.S. Fish and Wildlife Service.
58. McAuley DG, Keppie DM, Whiting Jr MR. 2020 American woodcock (*Scolopax minor*). In *Birds of the world* (ed. AF Poole), Ithaca, NY: Cornell Laboratory of Ornithology. See <https://birdsoftheworld.org/bow/species/amewoo/cur/introduction>.
59. Mazerolle MJ. 2020 Model selection and multimodel inference using the AICcmodavg package. See <https://cran.r-project.org/web/packages/AICcmodavg/index.html> (accessed 28 June 2023).
60. Burnham KP, Anderson DR. 2002 *Model selection and multimodel inference*. New York, NY: Springer.
61. Sheldon WG. 1967 *The book of the American woodcock*. Amherst, MA: University of Massachusetts Press.
62. Wingfield JC. 2005 Flexibility in annual cycles of birds: implications for endocrine control mechanisms. *J. Ornithol.* **146**, 291–304. (doi:10.1007/s10336-005-0002-z)
63. Maclean GL. 1972 Clutch size and evolution in the charadrii. *Auk* **89**, 299–324. (doi:10.2307/4084208)
64. Vander Haegen MW, Owen Jr RB, Krohn WB. 1994 Metabolic rate of American woodcock. *Wilson Bull.* **106**, 338–343.
65. Erckmann WJ. 1983 The evolution of polyandry in shorebirds: an evaluation of hypotheses. In *Social behaviour of female vertebrates* (ed SK Wasser), pp. 113–168. New York, NY: Academic Press.
66. MacLean Jr SF. 1969 Ecological determinants of species diversity of arctic sandpipers near Barrow, Alaska. Dissertation, University of California, Berkeley, CA, USA.
67. McAuley DG, Longcore JR, Sepik GF. 1990 Renesting by American woodcocks (*Scolopax minor*) in Maine. *Auk* **107**, 407–410. (doi:10.2307/4087628)
68. Dwyer TJ, Sepik GF, Dereleth EL, McAuley DG. 1988 *Demographic characteristics of a Maine woodcock population and effects of habitat management*. Fish and Wildlife Research 4. Washington, DC: U.S. Fish and Wildlife Service.
69. McAuley DG, Longcore JR, Sepik GF, Pendleton GW. 1996 Habitat characteristics of American woodcock nest sites on a managed area in Maine. *J. Wildl. Manag.* **60**, 138–148. (doi:10.2307/3802048)
70. Keppie DM, Redmond GW. 1988 A review of possible explanations for reverse size dimorphism of American woodcock. *Can. J. Zool.* **66**, 2390–2397. (doi:10.1139/z88-354)
71. Colwell MA. 2006 Egg-laying intervals in shorebirds. *Wader Study Group Bull.* **111**, 50–59.
72. Parris RW. 1983 Late nesting attempt by an American woodcock. *Kingbird* **33**, 253.
73. Whiting Jr MR, Bogus TG. 1982 Breeding biology of American woodcock in east Texas. In *U.S. Fish and wildlife service wildlife research report 14* (eds TJ Dwyer, GL Storm), pp. 132–138. Washington, DC: U.S. Fish and Wildlife Service.
74. Latham RM. 1956 Midwinter courtship flights of woodcock in Delaware. *J. Wildl. Manag.* **20**, 209. (doi:10.2307/3797437)
75. Vander Haegen WM. 1992 Bioenergetics of American woodcock during the breeding season on Moosehorn National Wildlife Refuge, Maine. Dissertation, University of Maine, Orono, ME, USA.
76. Wiley EN, Causey KM. 1987 Survival of American woodcock chicks in Alabama. *J. Wildl. Manag.* **43**, 414–420.
77. Slezak CR *et al.* 2024 Unconventional life history in a migratory shorebird: desegregating reproduction and migration. Figshare. (doi:10.6084/m9.figshare.c.7159242)